

H.O.P.E.: Suicide Prevention for American Indian and Alaska Native Crime Victims



H.O.P.E. saves lives

Acknowledgements

Several Education Development Center (EDC) staff members worked enthusiastically to realize our vision for an innovative, cross-cutting, and relevant curriculum that would help to equip crime victim advocates who serve American Indian/Alaska Native individuals with the information and skills they need to support their clients. The curriculum development team included Dr. Heidi Kar, Ms. Jennifer Myers, and Ms. Audrey Johnson. Special thanks are due to internal EDC reviewers, Dr. Jerry Reed, Ms. Terri Yellowhammer, Ms. Sadé Ali, Ms. Valda Grinbergs, and Mr. Adam Swanson and to the project's external board members, Dr. Jackie Campbell, Dr. Kate Cerulli, Dr. Joan Gillece, and Mr. William Kellibrew, for their feedback, guidance, and wisdom. We acknowledge our master trainers, Dr. Heidi Kar, Ms. Kayla Rae Wakeland, and Ms. Jennifer Myers, who skillfully conducted train-the-trainer workshops developing the skills of new H.O.P.E. trainers across the nation.

Access resources, videos, and additional information about the H.O.P.E. Suicide Prevention Training on the EDC website: <https://www.edc.org/hopeforadvocates>.

This product was supported by grant number 2017-VF-GX-K143, awarded by the Office for Victims of Crime, Office of Justice Programs, U.S. Department of Justice. The opinions, findings, and conclusions or recommendations expressed in this product are those of the contributors and do not necessarily represent the official position or policies of the U.S. Department of Justice.

Copyright © 2020 by Education Development Center.

Education Development Center (EDC) is a global nonprofit that advances lasting solutions to improve education, promote health, and expand economic opportunity. Since 1958, we have been a leader in designing, implementing, and evaluating powerful and innovative programs in more than 80 countries around the world. Visit [edc.org](https://www.edc.org).

Table of Contents

Module 1: Prepare Ourselves	3
Module 1 Overview	5
Welcome.....	6
Vignettes	9
Common Reactions	12
Managing Reactions	14
Conclusion	18
Module 2: Understand Suicide.....	19
Module 2 Overview.....	21
Using Data to Understand Suicide	22
Warning Signs	27
Risk Factors	31
Hopelessness.....	36
Connecting It All Together.....	37
Optional Guided Meditation.....	37
Conclusion	38
Module 3: Listen and Recognize	39
Module 3 Overview.....	41
Your Role as a Responder	42
Listen and Recognize	43
Vignettes	44
Common Trajectories.....	54
Conclusion	55
Module 4: Respond and Transition.....	57
Module 4 Overview	59
Introduction.....	60
Responding.....	60
Safer Suicide Triage.....	73
Conclusion	76
Module 5: Recharge Ourselves.....	77
Module 5 Overview.....	79
Introduction.....	80
Commitment to Self-Care	80

Vicarious Traumatization	83
Resources	85
Recharge Ourselves.....	86
Conclusion	90
Module 6: Consider the Complexities.....	91
Module 6 Overview	93
Introduction	94
Consider the Complexities.....	94
Unclear Situations	100
Conclusion	103
Module 7: Support the Family	105
Module 7 Overview.....	107
Introduction	108
Common Family Reactions	108
Supporting the Family	112
Conclusion	115
Module 8: Consider the System.....	117
Module 8 Overview	119
Introduction	120
Communities and Systems	120
Identifying Champions	123
Conclusion	124
Training Wrap-Up	124
References.....	127

Dear Advocates,

Education Development Center (EDC) is pleased to partner with the Office for Victims of Crime of the Department of Justice in presenting *H.O.P.E.: Suicide Training for American Indian and Alaska Native Crime Victims*. EDC's staff of licensed mental health providers, public health experts, crime victim advocates, and suicide prevention specialists, as well as our Board of Advisors, which includes experts from across crime victim advocate settings, including American Indian and Alaska Native settings, have contributed to the development of this curriculum.

This curriculum was designed to offer an effective, evidence-based, and context-specific suicide prevention training for advocates working across crime victim settings. Almost 100 trainers were trained to deliver the *H.O.P.E.: Suicide Prevention Training for Crime Victims* across the nation and they have, in turn, trained hundreds of advocates in this work so far.

This training manual has been informed by feedback from advocates who have been trained in the curriculum and who have in turn used their knowledge to train other advocates and to support their clients.

This manual is designed to be used to build advocates' own skills through self-guided study. We encourage any advocate—whether in domestic violence, child protection, or any other type of violence prevention advocacy role—to use these materials to increase their skills in suicide prevention among crime survivors.

If an organization would prefer to be trained by an established trainer in the H.O.P.E curriculum, please contact Dr. Kar at the email address below for information about trainers located in your geographic area.

Thank you for making a commitment to advancing your knowledge and skills in working with crime victims across American Indian and Alaska Native communities who struggle with suicidal thinking or behavior. Applying best practices and evidence-based tools and fine-tuning your suicide prevention skills will have a tremendous impact on those you serve as a crime victim advocate and on providing excellent care to the people you serve.

The U.S. Surgeon General's *2012 National Strategy for Suicide Prevention: Goals and Objectives for Action* rightly states that we all have a role to play in advancing suicide prevention and recommends specific ways we can all get involved. Some objectives urge collaborative engagement of individuals at risk, families, and concerned others. Other objectives describe the importance of addressing the most vulnerable groups. We know that crime victims are at increased risk for suicide, and as such, it is imperative that victim advocates be equipped with the best information, skills, and resources to enable them to support those under their care.

By using the *H.O.P.E.: Suicide Prevention Training for American Indian and Alaska Native Crime Victims* curriculum materials and learning how best to recognize and support crime victims at higher risk of suicide, you are helping to achieve the goals and vision of the National Strategy. The H.O.P.E. curriculum includes this self-guided learning manual, a project website with additional resources, and a shortened, online course version designed by the Office for Victims of Crime.

Since we know that it only takes one person to offer hope, your actions make a significant difference to help reduce the toll of suicide on our nation. Thank you for all you do to best support those who are vulnerable to suicide.

Heidi Kar, PhD, MHS
H.O.P.E. Co-Developer
Clinical Psychologist
Education Development Center
hkar@edc.org

Jennifer Myers, MA
H.O.P.E. Co-Developer
Mental Health Counselor
Education Development Center
jmyers@edc.org

Module 1: Prepare Ourselves



Look for Hints



Ask Openly about
suicide



Validate Pain



Explore reasons to live
and plan to stay safe

H.O.P.E. saves lives

Module 1 Overview

+ Topics Covered

- » Processing our own biases and reactions
- » Common reactions of a crime victim advocate
- » Recognition and awareness of linkages between our own reactions and the experiences and responses of those we serve
- » Strategies to manage and address biases and reactions

+ Learning Objectives

In Module 1, participants will:

- » Develop and improve skills in building self-awareness and increasing mindful processing of their own biases and reactions
- » Describe common reactions when working with crime victims at risk of suicide
- » Identify linkages between their own reactions and the experiences and responses of crime victims
- » Outline at least two strategies to manage their own reactions and to create a collaborative and non-adversarial stance

+ Materials

- » **Worksheet 1A: Vignettes**
- » **Worksheet 1B: Reflection Questions**
- » **Video: *Prevention Paradigm for Native Americans*** (5:16): Dr. Doreen Bird, Santo Domingo Pueblo, discusses challenges and opportunities in Native American suicide prevention: <https://www.sprc.org/video/native-americans>.

Welcome

Thank you for taking time away from your work to improve your suicide prevention knowledge and skills.

- » As you go through the materials, you will see the acronym H.O.P.E. referenced to assist you in remembering the steps you can take to instill hope and to help a victim through a suicidal crisis. H.O.P.E. stands for:

H.O.P.E. Saves Lives



Look for Hints



Ask Openly about suicide



Validate Pain



Explore reasons to live and plan to stay safe

- » We all have a role in offering hope, supporting, and advocating for those at risk for suicide.
- » We will be addressing several difficult topics throughout the training materials, including trauma, violence, victimization, and suicide risk.
- » While suicide may not be a word said often in some Native cultures and nations, suicide has likely affected many if not all of us. Often, we know someone who has thought about suicide or attempted suicide, or we have lost someone to suicide.
- » We are not alone in our experiences, and even though we have been through challenging times and loss, we can also heal and thrive. We have demonstrated this through centuries of healing, resilience, and growth in our nations and communities.
- » If you become uncomfortable, or are distressed at any point, please talk a support person, seek help, or reach out to one of the resources available to you in your community.
- » We have also included several suicide prevention resources. The first is the National Suicide Prevention Lifeline. It is a 24/7 talk line. You can also call this line and ask for help if you are concerned about a friend, person you are working with, or family member. Another resource is the Crisis Text Line, which can be used for any crisis. You may also have access to an Employee Assistance Program (EAP), local or regional crisis lines, and traditional healers.
 - Call the National Suicide Prevention Lifeline at 1-800-273-8255. This is a 24/7 talk line.

- Crisis Text Line – Text “Home” to 741-741. It is not just for someone thinking about suicide
- » Many of us don't spend this much time talking about suicide prevention. So, we encourage you to give yourself some extra care. Connect with colleagues, your loved ones, and the traditions that ground you.

Introduction

- » The training materials will cover eight modules:
 1. Prepare Ourselves
 2. Understand Suicide
 3. Listen and Recognize
 4. Respond and Transition
 5. Recharge Ourselves
 6. Consider the Complexities
 7. Support the Family
 8. Consider the System
- » Throughout these modules, keep in mind your role and the resources you have in your communities and apply what these materials address to your specific setting.
- » These self-guided training materials are designed to move suicide prevention beyond mental health. This training is intended to identify those who are at risk in their communities. Let's take a moment to hear from a suicide prevention expert about ways we can meet the challenges with connectedness and empowering our communities.
- » Play video – *Suicide Prevention in Native Communities*: <https://www.sprc.org/video/native-americans>.
- » Before we move on to Module 1, let's review the principles of a trauma-informed approach. These are foundational in our work. We encourage you to keep these in mind while reviewing the training materials.

Principles of a Trauma-Informed Approach

- » **Safety** – Staff and the people they serve feel physically and psychologically safe. Advocates consider safety as a core foundation to interactions, decisions, and responses.
- » **Trustworthiness and transparency** – Decisions are conducted with transparency, including the goal of building and maintaining trust among staff, clients, and family members of those receiving services.

- » **Peer support and mutual self-help** – These are integral and are understood as a key vehicle for building trust and establishing safety and empowerment.
- » **Collaboration and mutuality** – With this, there is a true partnering and leveling of power differences between staff. There is recognition that healing happens in relationships and in the meaningful sharing of power and decision-making.
- » **Empowerment, voice, and choice** – Individuals' strengths are recognized, built upon, and validated. Advocates aim to strengthen the experience of choice for clients, family members, and staff. This includes the awareness that every person's experience is unique and requires an individualized approach. A belief in resilience and in the ability of individuals, organizations, and communities to heal and recover from trauma is foundational. This is a strengths-based approach instead of a deficits-based approach.
- » **Cultural, historical, and gender issues** – Advocates actively move past cultural stereotypes and biases (e.g., based on race, ethnicity, sexual orientation, age, geography), offer gender responsive services, leverage the healing value of traditional cultural connections, and recognize and address historical trauma (Substance Abuse and Mental Health Services administration [SAMHSA], 2014).

For additional information – See SAMSHA's *Concept of Trauma and Guidance for a Trauma-Informed Approach* (2014) at https://store.samhsa.gov/product/SAMHSA-s-Concept-of-Trauma-and-Guidance-for-a-Trauma-Informed-Approach/SMA14-4884?referer=from_search_result

Medicine Wheel

- » Throughout the healing work that we do, we connect with the whole person, their whole being, and all that has come before in their lives and communities. The medicine wheel embodies the Four Directions: North, South, East, and West. It can also represent the dimensions of mental, emotional, spiritual, and physical. As we know, there are many meanings among our cultures of the medicine wheel. We bring this as a foundation to this training and our work honoring our health and healing currently and across generations.

Moving to Module 1: Prepare Ourselves

When working with individuals who have been the victims of crime, you will likely be working with individuals at risk for suicide. We bring our full presence to our work. This includes cultural beliefs about suicide, our attitudes, and our hopes. Before we cover what to do to help someone at risk for suicide, let's focus on ourselves and on our reactions. These reactions could be thoughts, emotions, sensations in our bodies, and body language. The word *suicide* itself is powerful and can evoke many different reactions. As we heard in the video, each nation may approach the topic and the word *suicide* differently.

Some of our personal responses may be useful and informative. Other personal reactions may be less helpful.

- » In this section, we will be addressing our own reactions, identifying some of our own biases, and sharing strategies for addressing these.

Vignettes

Vignettes and Reflection

- » Let's start by exploring some real-life scenarios. When reading and reflecting on these, focus on what comes up for you. Identify any thoughts, emotions, or physical sensations that you experience. Focus less on exactly what you would do and focus more on what prompts you to do that. For instance, if you want to get help right away, you might identify feeling worried or fearful or having a desire to fix it. If you think the person isn't really at risk for suicide, write about this.
- » Using **Worksheet 1A: Vignettes**, read a vignette and reflect on the questions. Reflect on your initial reactions without filtering them.

Worksheet 1A: Vignettes

- A. You are meeting with Harold, a 17-year-old member of an Indigenous nation. He reveals that his friend killed herself last week, the third person he has lost to suicide in the past two years. He tells you that he has a history of child sexual abuse from a family member. He says, "It just keeps happening. I don't think I can handle any more. It's too much—the suicides, the abuse. So many adults in our community are under the influence of alcohol and drugs and always seem angry or hopeless." When you ask him what strategies he is using to cope, he answers, "I don't know. Nothing, really. I just keep going through the motions, but I feel heavy and empty inside. Nobody will talk about it. Nobody lets you talk about the bad things, so they keep happening." You ask how it makes him feel that his community isn't discussing suicide or sexual abuse. He says, "Abandoned, lost. Sometimes I think, maybe they had the right idea—getting out."
- B. Danica is a 15-year-old Native American who identifies herself as two-spirit. She came to you through the Student Assistance Team at her school who referred her for unexplained absences, failing grades, and when she is in school, multiple trips to the nurse. You have been working with her for the past six months. She tells you that she is being followed home by groups of students who taunt her in school as well, and she feels unsafe both in school and at home. She recently deleted her Facebook page as a result of the many threatening comments about harm coming to her family. There is one particular student who seems to be obsessed with her and has been stalking her at home and contacting her on her cell phone numerous times a day. You have tried to engage her mom, but she works full time as medical assistant at a remote clinic on the reservation, and her drive is an hour each way. Danica's grandmother lives with them to care for Danica and her three younger brothers, but she speaks only her tribal language, and Danica is not fluent in it. Danica's dad is in the Army and is deployed in the Middle East. He has been away for nearly a year but makes periodic trips home. Danica feels as if she is a burden and has shamed her family. She has been telling you recently that, if she were not there, her family would be safer, and she feels as if she is a burden and shame to them.
- C. Raven is a 45-year-old Native American soldier in the National Guard. He served for 20 years in the Air Force and now lives on his ancestral lands of his reservation. He is recognized by his Chief and the Tribal Council as a leader in the community. He has always lived by Traditional Ancestral Ways, and he teaches language reclamation to the young people of the tribe. You first encounter Raven in the emergency room, where you provide advocacy services to him. He is being treated after violent sexual assault and attempted suicide immediately after the assault. Raven tells you that he has had recurring dreams of the lives that he and his troop took while he was on active duty, and his wife has taken to sleeping in another room following a night when he woke up screaming and throwing punches. He tells you that the spirits of those whose lives they took call to him. He cannot imagine returning back home to his wife and the tribal leaders if they know what happened to him. He has told the emergency department staff that he didn't mean to kill himself but tells you that he wishes he hadn't survived either the assault or the suicide attempt.
- D. Kateri is a 25-year-old Native American woman living on her reservation. For the past eight months, Kateri's twin sister, Angelica, has been missing from the home they share with their mother, father, and maternal grandmother. Kateri tells you that every day, she hears about missing and murdered Native women and girls, and she fears that her sister is among that number. You know from the investigation

Common Reactions

Introduction

You have reflected on several themes in our reactions to the scenarios.

Objectives

Participants will:

- » Describe common reactions when working with crime victims at risk of suicide
- » Identify linkages between their own reactions and the experiences and responses of crime victims

Key Points

- » Common reactions of the advocate can lead to over-control and under-responsiveness.
- » These are some common reactions that we may experience when working with a crime victim who is at risk for suicide:
 - **Fear** that we might say the wrong things, which might contribute to making things worse; that the person might die.
 - **Anxiety** about the unknown, about following procedures, about what to say or do, or about the impact that suicidal thinking or behaviors might have on the family or community.
 - **Anger**, especially if the person does not seem responsive to help.
 - **Helplessness** if it feels like you can't help or don't know what to do. This can occur if they are continuing to be victimized, and you feel helpless in how to protect them from it. This can also occur if there have been other suicide deaths in the community. You might feel as if you are helpless in your ability to stop clients from thinking of suicide.
 - **Hopelessness** that may occur when we feel less grounded, supported, and connected.
 - **Compassion** for the person we are working alongside.
 - **Empathy**, especially if we are able to understand their experiences and put ourselves in their shoes.
- » If we are feeling these emotions, it is possible that we might respond in one of two ways. We become over-controlling, or we withdraw. Neither of those responses are in the best interest of those individuals we are trying to serve.

- » Now, let's think about these reactions paired with some possible reactions of a victim. Possible reactions of the crime victim at risk can include keeping suicidal thoughts and behaviors secret, withdrawing, not showing up for or missing appointments, or not answering calls or checking in. Also, you might sense a change in rapport. This can feel like a power struggle at times. Consider that they may also be feeling many of the emotions we discussed the advocate experiencing. It is key for you in your role as an advocate to manage your own reactions to be able to fully support and respond. If you are well-versed with helpful, compassionate responses, it is possible for you to be more in sync with the individual. The person may be more likely to open up, disclose suicidal thoughts and behaviors, and seek support.
- » There are several common biases, or judgments, which a person might think about someone at risk for suicide. These include thinking the person is attention seeking, manipulative, and not serious enough (such as "They are only talking about it."), and that the person may be selfish. These judgments can cloud our reactions and are almost always not true about a person at risk for suicide. If the advocate thinks the person is attention seeking, it is important to remember that a lack of connectedness may be driving the person's suicidal thinking. Therefore, it may seem as if the person is seeking attention to meet this need. However, this is a real need that is causing pain and contributing to risk.
- » As advocates, we can balance and confront our judgments and biases with a genuine understanding of the experience of a person at risk for suicide. Of course, experiences are individual. Yet, research and experience has informed the field of suicide prevention.
- » It is likely the person in crisis is experiencing deep and unbearable pain and despair. Unresolved psychological pain can result in suicidal behavior (Levi-Belz et al., 2019; Shneidman, 1998). This pain can stem from unmet psychological needs. There are several possible factors which might be common for a people at risk:
 - **Thwarted belongingness/connectedness (Chu et al., 2017):**
 - Most people who think about suicide may not feel they are connected or belong.
 - **Perceived burdensomeness (Chu et al., 2017):**
 - Many persons at risk for suicide see themselves as a burden. While it may seem that they are being manipulative to get a response, they may be in crisis. It can also appear as if they are doing or saying things that are selfish, when they may believe their death relieves a burden rather than causes more pain.
 - **Ambivalence (Spencer-Thomas, 2017):**
 - Ambivalence is common for people in a suicidal crisis.
 - Like a teeter totter, a person's desire to live and desire to die can wax and wane. It can appear as if the person is not serious about dying, when what we know is that the person is likely very serious about ending their pain. They are struggling with how to do this in life. This balance can quickly tip towards the side of death without support and treatment.

- **Tunnel vision (Ellis, 2006; Oltmanns, 2017):**
 - One's thoughts can be so "tunneled" that they are only able to see the hurt, pain, and hopelessness. This is similar to a person who might see a dog and only experience this as a dog that might attack, even if all of the dog's signals and behaviors are showing that it is playful or wants affection.
- » For the crime victim, these suicidal thoughts can be triggered or intensified by the victimization or, what is for some, a re-victimization. The experience of the victim can include struggling to feel heard or believed and feeling blamed. This can be due to the nature of the victimization and the pervasiveness of the communication and actions by the offender blaming the victim. This can also be triggered as a result of immense stress that can be part of the criminal investigation process. Since the nature of a criminal investigation may also involve stress for the victim's family and community, concerns that they are a "burden" can be heightened.
- » **Reflection:** Thinking back on your work, can you think of a time that your own biases may have affected the care of a client or crime victim? Potentially, a situation where you may have over-controlled or under-responded to a person's crisis?

Managing Reactions

Introduction

There are many possible reactions to an individual that might be at risk for suicide. Some of these may be useful and assist in your work, while others may be hindrances and may contribute to you being out of sync with the individual.

There are several strategies we can employ to manage our reactions and moderate impacts on our work.

You may have developed and integrated your own strategies for managing your reactions in your work. Throughout this module, you might write your strategies down as you take notes. These strategies are not intended to be self-care strategies, although they may have a self-care benefit. Rather, these are to help us become aware of our reactions; see how they may be impacting our work, our colleagues, and our clients; and identify ways to address them.

Objectives

Participants will:

- » Outline at least two strategies to manage their own reactions and to create a collaborative and non-adversarial stance

Key Points

- » There are several strategies we can employ to manage our reactions and moderate impacts on our work.
- » In implementing these strategies, it helps if we have a culture of being able to support each other, especially when we see that a colleague may be over-responding, under-responding, or is not in sync with an individual. Some organizations have cultures that allow coworkers to hold each other accountable in a nonjudgmental way, such as being able to say, "I think this one hits really close to home, I wonder if I can help with it" Or, "It seems as if this person may be trying to tell you something. I wonder if there is something that might be blocking you from hearing it?"

Strategies to manage our reactions include:

- » **Self-reflection:**
 - Integrating self-reflection into our day-to-day work is key to identifying what is occurring in our reactions. Self-reflection can be done in many ways, such as through journaling, pausing, and informal processing with others. At times, we may have blind spots or might benefit from a formal consultation.
- » **Consultation:**
 - Consultation can be in the form of one-on-one consultation with a colleague, in a team or during staff meetings, or through other means, such as another agency.
 - Some organizations have a system through which a colleague or supervisor, typically a person with some expertise in working with a certain content area or population, can be accessed for consultation.
- » **Traditional and cultural practices:**
 - Traditional and cultural practices can be strengthening and grounding. It can be helpful to connect with your traditions and communities to know you are not doing this work alone and to connect with the deep Ancestral Knowledge of your nation.
- » **Spiritual practices:**
 - Spiritual practices can be very supportive in assisting to manage our own reactions, connect, and ground ourselves.
- » **Professional assistance:**
 - Professional assistance may be seeking therapy or medical treatment, if needed.
 - Our reactions often have roots. There will be times when we may not be able address a root or change the fruit developed through that root system without professional help. Working with clients who struggle with suicide can amplify our own trauma reactions. There is no shame in getting professional help, especially for the many of us who have been impacted by trauma, violence, and suicide.

» **Community and peer support:**

- Community and peer support can take lots of different forms. Peers can be our colleagues, peers who work in closely related professions, or peers who have similar experiences. Peers can play an important role in normalizing our reactions while also providing a safe space for growth.
- Community support can come from engaging in activities with each other, such as crafts, exercise, and more.

Reflection

- » Take time right now to reflect on ways you can best manage your own reactions by responding to the questions in **Worksheet 1B: Reflection Questions**.

- » In the work we do, language matters. It is important to use the words of the community we are working in and the language of the person we are working alongside. It is best to use terminology, such as *died by suicide* or *died of suicide*, as we would say for other medical issues. *Committed suicide* comes from terminology associated with committing a crime.
- » It is best to describe behavior directly rather than using phrases such as *an unsuccessful attempt*. We prefer to put the emphasis on life. Describe the actual behavior rather than labeling it as *manipulative*, *attention seeking*, or *a cry for help*. It is best to describe people using their qualities and character not as a diagnosis. The same is true for using terminology such as *we are dealing with a person who is suicidal*. Rather, *we collaborate with and support people who are in crisis*.
- » Quick tips to remember regarding language:
 - Died of/by suicide vs. ~~committed suicide~~
 - Suicide death/attempt vs. ~~successful/unsuccessful~~
 - Describe behavior vs. ~~manipulative/attention-seeking~~
 - Describe behavior vs. ~~suicide gesture/cry for help~~
 - Working with vs. ~~dealing with persons at risk for suicide~~ (Now Matters Now, 2018)


For more a digital image of language matters which can be shared with colleagues or posted in an office, see Now Matters Now at <https://www.nowmattersnow.org/wp-content/uploads/2018/10/LanguageMatters.pdf> The Now Matters Now offers tools, resources, and videos to apply Dialectical Behavior Therapy skills for managing suicidal crises and painful moments of life.

Conclusion

Key Points

- » It is common to have a variety of reactions to a person at risk for suicide:
 - Some of these reactions can be useful and inform our work, while other reactions could interfere with our relationship with an individual or impede our ability to be helpful.
 - Several strategies can be used to manage reactions. You don't have to do this work alone. A key point here is to become aware of your own reactions and use discernment regarding how they may be impacting you and your work.

Module 2: Understand Suicide



Look for Hints



Ask Openly about
suicide



Validate Pain



Explore reasons to live
and plan to stay safe

H.O.P.E. saves lives

Module 2 Overview

+ Topics Covered

- » Overview of key information about suicide
- » Review of the common signs of suicidal thoughts and behaviors
- » Review of risk factors associated with suicide

+ Learning Objectives

In Module 2, participants will:

- » Develop an awareness of the scope of suicidal thoughts and behaviors
- » Be able to name the signs of suicidal thoughts and behaviors across the life span
- » Be able to name the common risk factors for suicide
- » Identify hopelessness as a key risk factor for suicide

+ Materials

- » **Worksheet 2A: Look for Hints**
- » **Handout 2B: Warning Signs, Precipitating Factors, Risk Factors, and Protective Factors**
- » **Video: *Native Youth Are More Than Statistics* (13:37):** Ms. Elyssa Concha describes her own experience and that of her Native American people with suicide:
<https://www.youtube.com/watch?v=TMSyBmQt7iY>

Using Data to Understand Suicide

Introduction

In this section, there is a short review of data, then the content moves to warning signs and risk factors. Key take-home points include:

- » Data can help inform our work yet doesn't tell the full story.
- » The purpose of looking at data and understanding warning signs and risk factors is not to predict suicide. The purpose is to plan for response, treatment, and recovery.
- » Data are not representative of all persons and experiences.
- » It is key to be aware of warning signs.

Objectives


Participants will:

- » Develop an awareness of the scope of suicidal thoughts and behaviors at both the national and crime victim survivor level

Key Points

- » We have been focusing on our reactions, common experiences of the helper and the person in crisis, and ways to manage these reactions. While data and warning signs cannot predict who is at risk for suicide, this information can be useful to bring an understanding to suicide and help in planning our response.
- » We know that data are not just numbers on a page. The numbers cannot fully represent our people and our communities. We honor the people, communities, and the hurt that may be a part of our data. We also honor the resilience of our communities.
- » We know there are hundreds of tribal nations. As we heard in the video earlier, no data is representative of all nations and Native communities. Data varies in each community and is each community's story. The data we will discuss today are not intended to tell a specific nation's story or to even be generalized across all tribal nations or Native individuals.


Suicidal Behavior, United States 2018



www.sprc.org

Source: CDC, 2020; SAMHSA, 2019


- » This suicide behavior pyramid image is a good representation of opportunities to intervene Suicide Prevention Resource Center [SPRC], 2020b).
- » We may miss signs from individuals thinking about suicide and planning an attempt without the skills we will be focusing on in this guide.



Source: Centers for Disease Control and Prevention, 2020

- » Suicide disproportionately impacts non-Hispanic American Indian and Alaska Natives (Leavitt et al., 2018).
- » The suicide rate in the United States has increased by 33% since 1999. Yet, for Native communities the increase is even greater. The rate has increased 139% for Native women and 71% for Native men (Leavitt et al., 2018).
- » Data indicate what we all know: suicide deaths are a significant concern in our communities. Data show deaths in Indigenous populations occur more in the younger age groups. Youth aged 10–24 years accounted for 35.7% of AI/AN suicide deaths (9.8% aged 10–17 years, 25.9% aged 18–24 years). In comparison, 11.1% of suicides among Whites were in persons aged 10–24 years (2.5% aged 10–17 years, 8.6% aged 18–24 years). Almost 70 % of AI/AN decedents resided in nonmetropolitan areas, whereas most White decedents (over 70%) resided in metropolitan areas (Leavitt et al., 2018).
- » Men die by suicide more due to their tendency to use more lethal means (e.g., increased use of firearms), and there can be a lesser likelihood that they will access social support and/or treatment.


Suicide Rates for AI/AN Populations by Sex, United States 2009-2018



Source: Centers for Disease Control and Prevention, 2020

- » Suicidal behaviors and the risk of suicide are higher among the lesbian, gay, bisexual, transgender, transsexual, queer, questioning, and two-spirit (LGBTQ2S) population (We R Native, n.d.).
- » Often, when we think of suicide we don't think of younger children. Yet, young children can think about, attempt, and die by suicide. As we will discuss in detail later, abuse, trauma, and victimization can play a role in increasing the suicide risk. Native youth living on reservations are at a significantly higher risk for suicide and suicidal behavior, with lifetime rates of suicide and suicidal ideation much higher than the general population and Native youth in urban areas.

Suicide Rates for AI/AN Populations by Age, United States 2009-2018



www.sprc.org

*Non-Hispanic
Source: CDC, 2020

Source: Centers for Disease Control and Prevention, 2020

- » American Indian and Alaska Native women experience higher levels of violence than the general population. Almost 40% of Native women and 34% of Native men reported experiencing violence in the past year. This is more than 1 in 3 both men and women reporting they have experienced violence in the past year (Rosay, 2016).
- » There are many studies with data showing a connection between victimization and suicide risk or mental health concerns. Many of us know this and see this in our own work experiences and communities.
- » For clarity, let's pause and define terms. For this manual, a *victim* is defined as a person who suffers direct or threatened physical, emotional, or financial harm as a result of an act by someone else, which is a crime. This does not necessarily equate to trauma. *Trauma* relates more to the emotional and psychological response that can be related to victimization and other distressing experiences.
- » Let's remember that we are focusing on data regarding the intersection of victimization and suicide risk. Many people who experience victimization will experience recovery and healing, and they will thrive. Most will not think about, attempt, or die by suicide. Especially with trauma-informed care approaches and healing communities, survivors can be empowered and rebuild a sense of control.

- » We would be talking about statistics for this entire training if we included all of the research regarding crime victims and trauma and increased likelihood of mental disorders, symptoms, and substance use. This is just an overview. We can keep these statistics in mind as we look ahead to warning signs and risk factors for suicide.

Warning Signs

Introduction

In this section, we will first cover what warning signs are and then review the warning signs.

It can help to think of warning signs in two tiers. The first tier are warning signs of acute risk. These are signs that we have to act immediately. Think of these as an alarm, similar to a fire alarm. If the fire alarm went off right now, we would act immediately to get ourselves to safety.

The second tier are warning signs that might indicate a suicidal crisis yet need further exploration. If we use the analogy with fire, think of these signs as smelling smoke. If we smelled smoke, it could be coming from different sources: someone could have lit a candle, there could be a fire pit outside, or it could be indicative of the start of something concerning. We would need to explore further to find out. All warning signs need to be taken seriously and explored.

Objectives

Participants will:

- » Be able to name the main signs of suicidality across the life span

Key Points

H.O.P.E. Saves Lives



Look for Hints



Ask Openly about suicide



Validate Pain



Explore reasons to live and plan to stay safe

H.O.P.E. stands for:

- » Look for **Hints** means look for the warning signs that may be hints that a person is struggling.
- » Ask **Openly** about suicide – It is our role to go further into difficult topics.
- » Validate **Pain** – We need to meet people where they are at, in their pain, before we move into response and safety.
- » **Explore** reasons to live and plan to stay safe – Just like you do for planning for physical safety.

We will focus on looking for hints now. If we are not attentive to these warning signs, we might miss them. Look for hints.

- » Warning signs or hints of acute risk are signs of short-term or immediate risk. These hints need to be addressed for the safety of the victim as soon as possible.
- » Immediate risk – These hints need to be addressed for the safety of the victim as soon as possible:
 - Talking about wanting to kill oneself can include many different statements.
 - Some statements include “I’m going to kill myself” or “I would rather be dead and just might do something to get there” or “If such and such doesn’t happen, I will kill myself.”
 - Looking for a way to kill oneself, such as searching online or obtaining a gun.
 - Talking about feeling hopeless or having no reason to live (SPRC, n.d.).
- » These are additional hints that a victim may be at risk. These signs are like the smoke. You see them, and you need to look into them to determine if the person is at risk.
- » Warning signs of suicide risk include:
 - Talking about feeling trapped or in unbearable pain
 - Talking about being a burden to others
 - Increasing the use of alcohol or drugs
 - Acting anxious or agitated; behaving recklessly
 - Sleeping too little or too much
 - Withdrawing or feeling isolated
 - Showing rage or talking about seeking revenge
 - Displaying extreme mood swings (SPRC, n.d.; American Foundation for Suicide Prevention, 2020)

- » There are individual and cultural differences in the expression of distress and warning signs:
 - For instance, a military service member or a veteran may be less likely to directly disclose suicidal thinking. They may be more likely to discuss feeling trapped or burdened, increase substance use, and have changes in mood or sleep.
 - In some Native communities, talking about suicide could be an offense to spiritual traditions or to ancestors. Therefore, direct communication about suicide may be more difficult.
 - Certain spiritual beliefs and practices highly restrict the use of alcohol or drugs. For an individual who practices these beliefs, the use of alcohol or drugs even in a very small amount could be a serious warning sign.
 - For cultures that emphasize the group over the individual, the perception of being a burden to others or shaming the family could be a very serious sign of suicide risk.
- » It is important to consider the individual, their culture, and any dynamics that might impact their communication of warning signs. Imagine a crime victim who is gay and has a physical disability. Imagine if this person is used to hiding his sexuality and consistently working so others see him as capable, strong, confident, and independent. This person may not show what we might think of as warning signs for suicide risk. It is possible that hiding more, doing even more to show what is perceived as strength, and holding in any signs of difficulty or challenge may be signs of a struggle.
- » We encourage you to learn more about the groups and communities you are working with and keep this in mind during day-to-day interactions.

Video and Reflection

- » Using **Worksheet 2A: Look for Hints**, write down any warning signs, including specific quotes or statements, that would indicate risk you notice while watching the video. In parts of this video, the speaker will talk about statistics and her experiences. Write down the warning signs when she is describing her own experiences.
- » **Video: Native Youth Are More Than Statistics** (13:37): Ms. Elyssa Concha describes her own experience and that of her Native American people with suicide:
<https://www.youtube.com/watch?v=TMSyBmQt7iY>

Worksheet 2A: Look for Hints

Warning Signs for Suicide Risk

Suicidal ideation	
Increased substance use or substance misuse	
Sense of purposelessness	
Anxiety or agitation	
Feeling trapped	
Hopelessness	
Withdrawal or isolation	
Anger	
Recklessness	
Mood changes	



- » Reflect on the following question following the video:
 - What warning signs for suicide did you identify?
- » For many of us, this is a powerful video, and Ms. Concha poignantly shares her story. Thinking back to managing our reactions, if you were working with her, how would you manage your reactions to hear her hurt and identify her risk?
- » As was emphasized in the video and the title, our people are more than statistics, more than a number, and more than a set of warning signs.
- » We can be in tune with those in our communities to truly identify hints of suicide risk and support healing.

Risk Factors

Introduction

In addition to looking for warning signs to show hints of suicide risk, we can also be aware of risk factors, precipitating factors, and protective factors of suicide risk to help understand an individual's risk for suicide.

At the end of this section, **Handout 2B: Resources** will provide a general list of warning signs, precipitating factors, risk and protective factors, as well as risk and protective factors specific to Native populations.

Objectives

Participants will:

- » Be able to name the common risk factors for suicide

Key Points

- » Precipitating factors are events or situations in a person's life that can trigger a suicidal crisis. This list of precipitating factors from the Suicide Prevention Resource Center is summarized from research:
 - End of a relationship or serious relationship problems
 - Death of a loved one
 - Legal problems
 - Serious financial problems (SPRC 2020a)

- » You can see how a crime victim may have several of these factors, especially if the person has to engage a lawyer, move, or take additional steps for protection that could be costly, or if the person also has challenges working. These precipitating factors may be even more intense for someone with intersectional identities.
- » We can think of precipitating factors as things that might prompt or contribute to a suicidal crisis for a victim that is already coping with so much. Risk factors are things which might increase risk, but having risk factors does not mean a person is definitely going to attempt or even that they will ever think about suicide. Just as we know that among persons who have a family history of heart disease (a risk factor), most will never experience a heart attack, the same is true for risk factors for suicidal thoughts and/or behaviors.
- » **Reflection:** These precipitating factors are not Native specific. What are some precipitating events that might contribute to suicidal thinking in your tribal community?
- » Risk factors are characteristics of a person or their environment that increase the likelihood that they could die by suicide. This is similar to a family history of heart disease as a risk factor for a heart attack. Some of these risk factors, such as lack of access to health care or access to lethal means, may be modifiable. Often, risk factors are not modifiable, such as a prior suicide attempt. It is important to know these risk factors and to keep them in mind. We can think of risk factors like a threshold for concern. The more risk factors, the lower that threshold might be. As a result, we might use a more immediate response to connect that person with treatment and support even if they have denied any suicidal thinking or are showing only a couple of warning signs. Combine a recent victimization with risk factors and warning signs, and you can see how a person may be at significant risk. Risk does not equal prediction. Similar to suicide risk, there are many people at risk for a heart attack who will never experience a heart attack.
- » Risk factors include:
 - Prior suicide attempt(s)
 - History of abuse
 - Misuse of alcohol or other drugs
 - Mental disorders, particularly depression and other mood disorders
 - Access to lethal means
 - Knowing someone who died by suicide, particularly a family member
 - Social isolation
 - Chronic disease and disability
 - Lack of access to behavioral health care (American Foundation for Suicide Prevention, 2020; SPRC, 2020a)

- » Risk factors vary for individuals and groups. This training would be much longer if we reviewed all nations and all possible risk factors. These are additional risk factors for Native persons. Historical trauma, alienation, acculturation, and modern day trauma contribute to suicide risk in Native communities. These include a history of interpersonal violence, barriers to help-seeking, social disintegration, substance misuse, discrimination, isolation, disconnection with traditional ways, suicide clusters, and economic instability. As with the warning signs earlier, it is important to know any particular risk factors in the communities in which you are working.
- » **Reflection:** What are risk factors for suicide that are specific to your community?
- » Protective factors are characteristics that might keep a person from thinking about or attempting suicide. Protective factors include:
 - Being a parent
 - Effective behavioral health care
 - Connectedness to individuals, family, community, and social institutions
 - Life skills (including problem-solving skills and coping skills, ability to adapt to change)
 - Self-esteem
 - Sense of purpose or meaning in life
 - Cultural, religious, or personal beliefs that discourage suicide (SPRC, 2020a)
- » For crime victims, children may serve as a risk or a protective factor. Consider a victim that firmly believes her children need her and, therefore, would not act on suicidal thoughts. She may believe she does not want to add more trauma to her children's lives and wants to live for them. Now, consider if an abusive ex-partner talks about or tries to hurt the children to get back at the victim. In this situation, the victim may be more likely to think about or even attempt suicide as a way to protect the children from harm.
- » Knowing the risk factors for suicide, it might make sense that some of these protective factors are opposites of those risk factors.
- » A key strategy in assisting a person at risk for suicide is to increase protective factors and decrease risk factors. This is similar to increasing coping and life skills and decreasing unhealthy coping responses that might display as warning signs.

Handout 2B: Warning Signs, Precipitating Factors, Risk Factors, and Protective Factors

Warning Signs

Immediate Risk

The following three behaviors indicate that immediate action and support is needed through a crisis response or by contacting a mental health professional:

- » Talking about wanting to die or to kill oneself
- » Looking for a way to kill oneself, such as searching online or obtaining a gun
- » Talking about feeling hopeless or having no reason to live

Serious Risk

Other behaviors may also indicate a serious risk—especially if the behavior is new, has increased, and/or seems related to a painful or traumatic event, loss, or change:

- » Talking about feeling trapped or in unbearable pain
- » Talking about being a burden to others
- » Increasing the use of alcohol or drugs
- » Acting anxious or agitated; behaving recklessly
- » Sleeping too little or too much
- » Withdrawing or feeling isolated
- » Showing rage or talking about seeking revenge
- » Displaying extreme mood swings

Precipitating Factors

Precipitating factors are stressful events that can trigger a suicidal crisis in a vulnerable person. Examples include:

- » End of a relationship or marriage
- » Death of a loved one, especially if by suicide
- » An arrest
- » Serious financial problems

Risk Factors

Risk factors are characteristics of a person or their environment that increase the likelihood that they will die by suicide (i.e., suicide risk).

Major risk factors for suicide include the following:

- » Prior suicide attempt(s)
- » Misuse of alcohol or drugs
- » Mental disorders, particularly depression and other mood disorders
- » Access to lethal means
- » Knowing someone who died by suicide, particularly a family member/tribal member
- » Social isolation
- » Chronic disease and disability
- » Lack of access to behavioral health care

Risk Factors Specific to Native Communities and Individuals

Risk factors can vary by nation, age, culture, sex, and other characteristics.

Risk factors that can be more specific to Native communities and individuals include a history of interpersonal violence, barriers to help-seeking, social disintegration, substance misuse, discrimination, isolation, disconnection with traditional ways, suicide clusters, and economic instability.

Protective Factors

Protective factors are personal or environmental characteristics that help protect people from suicide.

Major protective factors for suicide include the following:

- » Strong family connections
- » Strong connections to one's nation and traditional ways
- » Effective behavioral health care
- » Connectedness to individuals, family, community, and social institutions
- » Life skills (including problem-solving skills and coping skills, ability to adapt to change)
- » Self-esteem and a sense of purpose or meaning in life
- » Cultural, religious, or personal beliefs that discourage suicide



Hopelessness

Introduction

- » Hope can be a significant factor in contributing to a crime victim's will to live. It is key to identify hopefulness and hopelessness.
- » This section will focus on the importance of drawing out a person's level of hope, paying attention to factors which might contribute to changes in hopelessness, and ways to ask about hopelessness.
- » It is important to remember that hope is more than future orientation. A person can have plans or thoughts about the future, but this does not mean that they have hope about those plans.

Objectives

Participants will:

- » Be able to identify hopelessness as a key risk factor for suicide

Activity

- » Let's brainstorm ways to identify hope, hopelessness, and hopefulness.
- » **Reflection:**
 - What are some questions you can ask to identify levels of hope? Hopelessness? Hopefulness?
 - What are behaviors and nonverbal communications that you look for to know if a person is feeling hopeless?
- » Social isolation often goes hand in hand with hopelessness. This could include a victim that moved from a reservation to an urban area or, like we watched in the video, an adolescent feeling isolated as one of the only Native persons at her school.
- » These are some ideas for identifying the intensity of hopelessness. It is best if you ask these questions once a relationship has been established:
 - On a scale of 1–10, with 10 being so hopeless it feels like you can't go on, where would you say you are right now?
 - When was the last time you felt like you were closer to a 1 on this scale?
 - What do you want to happen that would feel like there would be some hope?

Connecting It All Together

Introduction

We have reviewed data related to suicide and the intersection with crime victims, warning signs, risk factors, protective factors, precipitating events, and hopelessness. Now, let's bring it all together.

Key Points

- » As with violence prevention, one cannot predict when and how someone might attempt or die by suicide. All of these factors come together in our planning and in our response. The purpose of learning about suicide prevention is not prediction but planning.
- » Since triggering events can be sudden, it is important to look for hints and plan for safety.
- » A study conducted by several researchers at Harvard found that the median onset for certainty about the place to attempt a suicide was 30 minutes prior to the attempt, and the median time for the final decision to attempt a suicide was 5 minutes before the attempt (Millner et al., 2016). We may not be able to predict when someone might act on suicidal thoughts. Yet, we can increase our awareness to see the hints, mutually work to develop safety plans, and improve support that can be available at a time of crisis.

Optional Guided Meditation

Introduction

This is an optional guided meditation (approximately 3–5 minutes) that can offer a break from the intensity of the content. Feel free to use an alternative meditation, if you wish.

Let us take a few minutes, now, to step away from the material we've been discussing and give ourselves a mental break. Read the text below and take a few minutes to engage in a brief meditation exercise called "Leaves on a Stream."

“Leaves on a Stream” Exercise (Harris, 2009)

1. Sit in a comfortable position and either close your eyes or rest them gently on a fixed spot in the room.
2. Visualize yourself sitting beside a gently flowing stream with leaves floating along the surface of the water. Pause 10 seconds.
3. For the next few minutes, take each thought that enters your mind and place it on a leaf... let it float by. Do this with each thought—pleasurable, painful, or neutral. Even if you have joyous or enthusiastic thoughts, place them on a leaf and let them float by.
4. If your thoughts momentarily stop, continue to watch the stream. Sooner or later, your thoughts will start up again. Pause 20 seconds.
5. Allow the stream to flow at its own pace. Do not try to speed it up and rush your thoughts along. You are not trying to rush the leaves along or “get rid” of your thoughts. You are allowing them to come and go at their own pace.
6. If your mind says, “This is dumb,” “I’m bored,” or “I’m not doing this right,” place those thoughts on leaves, too, and let them pass. Pause 20 seconds.
7. If a leaf gets stuck, allow it to hang around until it’s ready to float by.
8. If the thought comes up again, watch it float by another time. Pause 20 seconds.
9. From time to time, your thoughts may hook you and distract you from being fully present in this exercise. This is normal. As soon as you realize that you have become sidetracked, gently bring your attention back to the visualization exercise.

Conclusion

Key Points

- » Preventing suicide as an advocate requires an understanding of suicide; looking for hints of suicide risk; and recognizing warning signs, risk factors, precipitating factors.
- » Understanding these factors on a population level enables advocates to look for them in their communities and the individuals they are supporting.
- » In the next section, we will discuss the scope of your role. Your role is a recovery-oriented, strengths-based one. With an understanding of the suicidal crisis and the impact of trauma, typically, there is not one line we can say to someone who is struggling that will pull them out of a suicidal crisis, although we may wish this were the case. In the next section, we will discuss an approach to listening and responding which is helpful to those who struggle with suicidal thoughts and behaviors.

Module 3: Listen and Recognize



Look for Hints



Ask Openly about
suicide



Validate Pain



Explore reasons to live
and plan to stay safe

H.O.P.E. saves lives

Module 3 Overview

+ Topics Covered

- » How and what to listen for regarding suicidal thoughts and feelings
- » Asking openly about suicide using skilled questions
- » Developing the ability to respond with both indirect and direct methods to explore suicide risk
- » Common trajectories to suicide among crime victims

+ Learning Objectives

In Module 3, participants will:

- » Recognize the scope of their role when working with an individual at risk for suicide
- » Be able to identify subtle references to feelings of hopelessness and burdensomeness
- » Be able to effectively elicit key warning signs of suicide
- » Develop confidence to respond with both indirect and direct methods to explore suicide risk

+ Materials

- » **Handout 3A: Vignettes**
- » **Handout 3B: Columbia-Suicide Severity Rating Scale**
- » **Handout 3C: Role-Plays**

Your Role as a Responder

Introduction

- » This section moves from a focus on what to look for and begins to add how to actively listen, identify, and respond. We will focus on identifying and eliciting the hints we outlined in the last section.
- » Active and intentional listening for warning signs and using skilled questions are key to our role of identifying if a victim is at risk. It is especially important to use active listening and to match the victim in their style and manner of communication when someone has experienced a traumatic event, like a violent crime. There can be many barriers to reporting a crime for some individuals, groups, and communities, such as minority, immigrant, Native, and LGBTQ2S communities.
- » We will outline what to listen for, how to ask skilled questions, and how to respond so as to potentially interrupt a path toward suicidal behaviors.

Objectives

Participants will:

- » Recognize the scope of their role when working with an individual at risk for suicide
- » Be able to identify subtle references to feelings of hopelessness and burdensomeness
- » Be able to effectively elicit key warning signs of suicide
- » Develop confidence to respond with both indirect and direct methods to explore suicide risk

Key Points

- » As we go through this section, it is important to remember our roles and to have clear expectations of the advocate's core responsibility with suicide prevention: to identify risk, not to assess and treat.
 - If someone seems to be in need of CPR, what is the first thing you do?
 - If someone seems to be in need of CPR, what is the first thing you do?
 - Then, what are a few next steps?
 - Let's think of this situation as similar to the fire alarm example that we discussed earlier. A person who might be in cardiac arrest is the highest level of concern, just like a fire alarm going off or seeing a fire.
 - What about a person who is complaining of shortness of breath? What else would you want to know?

- » It is your role as a trained responder to identify the hints and ask more questions to determine the appropriate response, then to initiate this response. It is not your role to treat a heart attack, install a stent, and perform cardiac rehab. Nor is it your role to determine the level of risk. Is it really a heart attack? A panic attack? Once you see the signs, the assessment by a trained medical professional is urgently indicated. Your role is to identify, support, and help a person remain safe and connect to treatment. Similarly, in victims' services, your role is supportive and empowering with an emphasis on safety.

Listen and Recognize

Objectives

Participants will:

- » Be able to identify subtle references to feelings of hopelessness and burdensomeness
- » Be able to effectively identify key warning signs of suicide
- » Develop the ability to respond with both indirect and direct methods to explore suicide risk

Key Points

- » Review and practice looking for hints.

H.O.P.E. Saves Lives



Look for Hints



Ask Openly about suicide



Validate Pain



Explore reasons to live and plan to stay safe

Vignettes

- » Please read over a vignette in **Worksheet 3A: Vignettes** using a different vignette than you read in Module 1. Make note of hints of suicide risk you identify in the vignette and pay particular attention to recognizing warning signs.
- » Write down anything that may provide a hint someone is at risk for suicide, such as:
 - Warning signs
 - Risk factors
 - Precipitating factors
 - Hopelessness
 - Burdensomeness
 - Purposelessness

Worksheet 3A: Vignettes

- A. You are meeting with Harold, a 17-year-old member of an Indigenous nation. He reveals that his friend killed herself last week; the third person he has lost to suicide in the past two years. He tells you that he has a history of child sexual abuse from a family member. He says, "It just keeps happening. I don't think I can handle any more. It's too much—the suicides, the abuse. So many adults in our community are under the influence of alcohol and drugs and always seem angry or hopeless." When you ask him what strategies he is using to cope, he answers, "I don't know. Nothing, really. I just keep going through the motions, but I feel heavy and empty inside. Nobody will talk about it. Nobody lets you talk about the bad things, so they keep happening." You ask how it makes him feel that his community isn't discussing suicide or sexual abuse. He says, "Abandoned, lost. Sometimes I think, maybe they had the right idea—getting out."
- B. Danica is a 15-year-old Native American who identifies herself as two-spirit. She came to you through the Student Assistance Team at her school, who referred her for unexplained absences, failing grades, and, when she is in school, multiple trips to the nurse. You have been working with her for the past six months. She tells you that she is being followed home by groups of students who taunt her in school as well, and she feels unsafe both in school and at home. She recently deleted her Facebook page as a result of the many threatening comments about harm coming to her family. There is one particular student who seems to be obsessed with her and has been stalking her at home, contacting her on her cell phone numerous times a day. You have tried to engage her mom, but she works full time as a medical assistant at a remote clinic on the reservation, and her drive is an hour each way. Danica's grandmother lives with them to care for Danica and her three younger brothers, but she speaks only her tribal language, and Danica is not fluent in it. Danica's dad is in the Army and is deployed in the Middle East. He has been away for nearly a year but makes periodic trips home. Danica feels as if she is a burden and has shamed her family. She has been telling you recently that, if she were not there, her family would be safer, and she feels as if she is a burden and shame to them.
- C. Raven is a 45-year-old Native American soldier in the National Guard. He served for 20 years in the Air Force and now lives on his ancestral lands of his reservation. He is recognized by his Chief and the Tribal Council as a leader in the community. He has always lived by Traditional Ancestral Ways, and he teaches language reclamation to the young people of the tribe. You first encounter Raven in the emergency room, where you provide advocacy services to him. He is being treated after a violent sexual assault and attempted suicide immediately after the assault. Raven tells you that he has had recurring dreams of the lives that he and his troop took while he was on active duty, and his wife has taken to sleeping in another room following a night when he woke up screaming and throwing punches. He tells you that the spirits of those whose lives they took call to him. He cannot imagine returning back home to his wife and the tribal leaders if they know what happened to him. He has told the emergency department staff that he didn't mean to kill himself but tells you that he wishes he hadn't survived either the assault or the suicide attempt.
- D. Kateri is a 25-year-old Native American woman living on her reservation. For the past eight months, Kateri's twin sister, Angelica, has been missing from the home they share with their mother, father, and maternal grandmother. Kateri tells you that every day, she hears about missing and murdered Native

Key Points

- » As we look for hints, we need to **ask openly**, in a sensitive manner, about suicide to determine if a person is at risk for suicide. This helps us get more information so we know how to support the victim. By doing so, we can start to offer hope, and we know that hope saves lives.

H.O.P.E. Saves Lives



Look for Hints



Ask Openly about suicide



Validate Pain



Explore reasons to live and plan to stay safe

- » Indirect questions focus on understanding the context of factors that are closely linked to suicide for most people, including whether they feel hopeless, feel as though they are a burden to others, think their loved ones would be better off without them, etc.
- » Examples of indirect questions:
 - Have you wished you were dead?
 - Have you wished you could go to sleep and not wake up?
 - Do you wish you could go away and never come back?
 - Have you had thoughts of harming yourself?
 - Do you feel like you are a burden to others?
- » Direct questions specifically ask about the person's wish to die or plans to end their lives.
- » Each of these questions will likely result in unique responses, giving you different information. For instance, a positive answer to "Do you feel like you are a burden to others?" will provide you with a significant sign. Yet, it doesn't directly ask the person if they wish they were dead.
- » It is important to know if a person is thinking of harming themselves.
- » Consider the difference between asking if a person has had thoughts of harming themselves and asking directly about suicide.
 - A person might say no to harming themselves but yes to more direct questions about suicide.

Activity

Identify your go-to questions for asking directly and yet sensitively about suicide. Take about 5–10 minutes to write out the ways you ask both directly and indirectly about suicidal thinking, behaviors, and plans.

- » In addition to your go-to questions, another way to elicit accurate information is to ask, “Have you felt so _____ (depressed, hopeless, ashamed, much pain) that you have thought about suicide?” Asking in this way can help to normalize these struggles. It can make it more likely for the person to give a genuine answer.
- » Another technique is gentle assumption. By asking when the last time was the person thought about suicide, you assume they have thought about suicide. This might make it easier for the person to share suicidal thoughts.
- » A tool that can assist in asking about suicide is the Columbia-Suicide Severity Rating Scale (C-SSRS). The Columbia-Suicide Severity Rating Scale is a standard, validated tool using plain-language questions that anyone can ask. A copy of the C-SSRS is included in [Handout 3B](#).
- » There are many tribal and Indian Health Service systems that use the Columbia-Suicide Severity Rating Scale. This is a tool that can be used in your communities. Yet, asking openly about suicide should not be dependent on one tool or one standard scale. As we know, people and communities are not the same. It is important to ask openly using our advocacy skills, building a relationship, and applying the questions we just reviewed. Let’s review this additional resource that we could add to our skilled connecting.

Handout 3B: Columbia-Suicide Severity Rating Scale – Screen Version, Recent

SUICIDE IDEATION DEFINITIONS AND PROMPTS	Past month	
	YES	NO
Ask questions that are bolded and <u>underlined</u> .		
Ask questions 1 and 2		
1. <u>Have you wished you were dead or wished you could go to sleep and not wake up?</u>		
2. <u>Have you had any actual thoughts of killing yourself?</u>		
If YES to 2, ask questions 3, 4, 5, and 6. If NO to 2, go directly to question 6.		
3. <u>Have you been thinking about how you might do this?</u> e.g. "I thought about taking an overdose but I never made a specific plan as to when where or how I would actually do it...and I would never go through with it."		
4. <u>Have you had these thoughts and had some intention of acting on them?</u> As opposed to "I have the thoughts but I definitely will not do anything about them."		
5. <u>Have you started to work out or worked out the details of how to kill yourself? Do you intend to carry out this plan?</u>		
6. <u>Have you ever done anything, started to do anything, or prepared to do anything to end your life?</u> Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump, or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.		
If YES, ask: Was this within the past three months?		

- Low Risk
- Moderate Risk
- High Risk

Source: Columbia Lighthouse Project, 2016



- » [The Columbia Lighthouse Project website](#) has resources, research, toolkits, and videos to assist with the use of the C-SSRS.
- » The C-SSRS captures both suicidal thoughts and suicidal behaviors, allows a quicker crisis response, and reduces the burden of unnecessary steps. Using a standard screening tool enables clearer terminology across advocates and systems. There are multiple resources with the C-SSRS. We will be focusing on the Screener – Recent.
- » The C-SSRS defines terms related to suicide risk. As with all of our work, it is essential to consider culture and context with language. For instance, the Native communities we are working in may use other terminology for terms related to suicide and suicidal behavior. We will review the terms used the Columbia-Suicide Severity Rating Scale. I encourage you to think of the terms you would use with the people you are working with, in their language.
- » Let us review the terms as they are defined in the C-SSRS:

Suicidal ideation: Thinking about killing oneself (Columbia Lighthouse Project, 2016).

Suicide attempt: A potentially self-injurious act committed with at least some wish to die as a result of the act. Behavior was in part thought of as a method to kill oneself. Intent does not have to be 100%. If there is any intent/desire to die associated with the act, then it can be considered a suicide attempt. There does not have to be any injury or harm, just the potential for injury or harm. If a person pulls the trigger while the gun is in their mouth, but the gun is broken so no injury results, this is still considered an attempt (Columbia Lighthouse Project, 2016).

- **Inferring intent:** Even if an individual denies the intent/wish to die, it may be inferred from the behavior or circumstances; for example, a highly lethal act that is clearly not an accident so no other intent but suicide can be inferred (e.g., a gunshot to the head, jumping from the window of a high story in a building). Also, if someone denies intent to die, but they thought that what they did could be lethal, intent may be inferred (Columbia Lighthouse Project, 2016).
- **Interrupted attempt:** When the person is interrupted (by an outside circumstance) from starting the potentially self-injurious act (if not for the interruption, an actual attempt would have occurred). This includes when a person has pills in hand but is stopped from ingesting. Once they ingest any pills, the act becomes an attempt rather than an interrupted attempt. However, if a person is poised to jump, but is grabbed and taken down from ledge, the act is an interrupted attempt (Columbia Lighthouse Project, 2016).
- **Aborted or self-interrupted attempt:** When a person begins to take steps toward making a suicide attempt but stops themselves before they actually have engaged in any self-destructive behavior. Examples are similar to interrupted attempts, except that the individual stops themselves, instead of being stopped by

something or someone else (Columbia Lighthouse Project, 2016).

Preparatory acts or behavior: Acts or preparation towards imminently making a suicide attempt. This can include anything beyond a verbalization or thought, such as obtaining a specific method or preparing for one's death by suicide such as writing a suicide note (Columbia Lighthouse Project, 2016).

- » When you look for hints and see the need to ask openly about suicide, use the following questions excerpted from the C-SSRS to help you ask directly (Columbia Lighthouse Project, 2016):
 - First, you can ask questions 1 and 2.
 1. Have you wished you were dead or wished you could go to sleep and not wake up?
 2. Have you had any actual thoughts of killing yourself?
 - If YES to question 2, ask questions 3, 4, 5, and 6. If NO to 2, go directly to question 6.
 3. Have you been thinking about how you might do this?
 4. Have you had these thoughts and had some intention of acting on them?
 5. Have you started to work out or worked out the details of how to kill yourself? Do you intend to carry out this plan?
 6. Have you ever done anything, started to do anything, or prepared to do anything to end your life?
 - Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump, or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.
 - If yes, ask: Was this within the past 3 months?
- » Notice, we are asking about suicidal ideation and suicidal behavior with all persons. What we know is that sometimes people in crisis might say they did not want to kill themselves or might deny suicidal thinking, yet they have engaged in some suicidal behavior in the recent past. This tool can be useful to ask about both ideation and behavior, which is key information when supporting a crime victim.
- » When using this scale and terminology, it is important to consider age, developmental level, culture, and any language barriers. Overall, these questions are good to use with persons aged 12 and up.
- » Consider your concerns in asking directly about suicide. What would prevent you from

doing this?

- » Common responses include the following:
 - **Fear** that they may make the person think about suicide:
 - Research shows that people at risk feel relief that they have been asked directly, and asking directly does not increase the likelihood a person will think about suicide.
 - **What if they say yes:**
 - Remember your role. You are there to listen, identify, assist with safety, and connect them to resources. You don't have to do this alone. We will talk more about how to respond throughout the training. First, you might take a moment to center yourself, such as by taking a deep breath. Acknowledge and support their pain. Validate their pain and connect with the part of them that wants to live. Explore reasons to live and connect them to treatment.
 - **What if they don't tell the truth:**
 - It's okay to trust your gut. If you think they may still be at risk, seek consultation, collaboratively establish a safety plan, and engage in treatment.
- » Now, we are going to practice asking directly about suicide using example scenarios in **Handout 3C: Role-Plays**. If possible, work with a colleague to role-play the scenarios provided in the text below. If it is not possible to practice right now, set an intention to practice when you are able. Take time now to read through one of the scenarios and write out how you would ask directly about suicide.
- » Take about 5 minutes for each of you to role-play. Each person should ask openly about suicide in this scenario. If you are role-playing a crime victim, you can respond to being asked about suicide in any way you feel works for you.
- » After you practice this exercise, discuss what went well and any difficulties or challenges. It is important to get to direct questions to ensure you understand the level of risk facing the person that you are working with.



Handout 3C: Role-Plays

The person role-playing the crime victim: Read the scenario to yourself. You can give the person acting as the advocate a little bit of background such as your age and what recently happened. Then, say the statement at the end of the scenario to start the role-play.

Scenario 1

You are a 24-year-old who has recently experienced physical and sexual assault. You have experienced trauma in your childhood home in the past, have often used substances to cope, and feel as if you have very limited supports. It seems best for you to move for your safety and mental health, but you feel like you don't have any options, and you don't want to move back to your childhood home. You have started to use substances more often to cope with the symptoms of the acute stress response and the many triggers you experience throughout the day. You tell the advocate, "I just don't feel like I have any options. I don't know how to recover from this, and I am not sure it is even worth me trying. Maybe it would be best if I just didn't even bother and waste your time. I just feel like giving up."

Scenario 2

You are a 72-year-old who has recently experienced elder abuse by your child, and you have connected with the victim advocate. You feel like a failure, as if you haven't raised your child well and have failed at helping him to be successful in life, especially regarding cultural beliefs about respecting and honoring elders. You blame yourself for the recent violence. You feel guilty, ashamed, helpless, and hopeless. While you have experienced the support of your nation, family, and friends in the past, you don't feel you can reach out to any of them or talk to them about this because you feel too ashamed. You don't see the situation as improving, don't feel you can cut off your child from your life, and don't think the advocate can help you. You state "Things can't get any worse. I think I would be better off dead."

Common Trajectories

Introduction

In addition to understanding data, signs, factors, and responding, it is helpful to understand some common trajectories in crime victim populations. We will review common paths to suicidal behavior. As with most things in our field, individual variations can occur, and these paths are not representative of every possible trajectory.


Objectives

Participants will:

- » Be able to identify common trajectories to suicide in crime victim populations

Key Points

- » Several factors can be common in a person's path toward a suicide attempt.
- » Adverse childhood experiences can affect biological factors, psychiatric and health functioning, and psychosocial development across the lifespan. The interaction of these factors with later-life stressors can lead to suicidal thoughts and behaviors in later life. There are multiple pathways towards suicidal thoughts and behaviors. Yet, we know that trauma, including the varied impacts of ACEs persist, thus, effecting suicide risk throughout the lifespan and into late life (Sachs-Ericsson et al, 2015).




- » We referenced this earlier. We cannot predict the if, when, and how of a suicide attempt. But we can respond in a timely manner to support a person in crisis and help prevent an attempt. When thinking about responding, it is not your role to judge or assess if a person might attempt. Keep in mind that suicidal thinking can occur for years before an attempt, or it can occur the week and even hours before an attempt. Suicide risk assessment is key to determining if someone is at risk. Your role is to identify, support, and connect to a mental health professional for this assessment and treatment.

Conclusion

Key Points

- » It is essential to recognize hints, ask openly about suicide, and stay within the scope of your role.

Module 4: Respond and Transition



Look for Hints



Ask Openly about
suicide



Validate Pain



Explore reasons to live
and plan to stay safe

H.O.P.E. saves lives

Module 4 Overview

+ Topics Covered

- » Safer suicide triage
- » Responding to a person at risk for suicide

+ Learning Objectives

In Module 4, participants will:

- » Build confidence to respond to persons at risk and connect with resources
- » Increase awareness of the types of practice-based interventions available for suicide to be able to describe options to victims.
- » Develop a framework for supporting a person at risk for suicide in receiving the next level of care.

+ Materials

- » **Handout 4A: Columbia-Suicide Severity Rating Scale – Screen with Triage Points for Victim Advocates**
- » **Handout 4B: Resources**
- » **Handout 4C: Patient Safety Plan Template**
- » **Worksheet 4D: Mental Health Supports**
- » **Video: *Suicide survivor on the best part of being alive? 'Everything'* (3:40):**
<https://www.usatoday.com/videos/news/2018/11/20/suicide-survivor-profile-shelby-rowe/1737400002/>

Introduction

Introduction

- » This section covers the response needed from the advocate after a person has been identified at risk.
- » Suicide safer triage is a step-by-step guidance on how to decide when a person is at risk and needs to be referred to a mental health provider for assessment.
- » Now that we have discussed managing our own reactions, hints to look for, and how to ask openly about suicide, let's talk about ways to respond when working with a person that you have identified as at risk.
- » We will address key skills and strategies in responding. Your specific role and resources in your system may allow you to do some of these things as a team. We encourage you to consider what you would do in your role and think about how you would respond to suicide risk applying the information covered in this section.

Key Points

- » Develop confidence to respond to the persons at risk and connect with resources.
- » Develop awareness of the types of evidence-based interventions available for suicide to be able to describe options to clients.
- » Develop a framework for transitioning survivors to the next level of care.

Responding

Introduction

In this section, we will be discussing the advocate's response and suicide safer triage.

Objectives

Participants will:

- » Build confidence to respond to persons at risk and connect them with resources

Key Points

- » Understanding the levels of risk and taking actions based on them is important.
- » For persons who might be lower risk—remember H.O.P.E. saves lives:

H.O.P.E. Saves Lives



H–Look for **H**ints

- If you identify hints, then



O–Ask **O**penly about suicide

- If you identify suicide risk, refer for mental health assessment



P–Validate **P**ain

- With the person at risk



Explore reasons to live and **E**stablish a safety plan

- Support safety
- Follow up
- Engage them in care

- » In addition to asking openly about suicide, validating pain, exploring reasons to live, and planning to stay safe, assist the person at risk to engage in mental health assessment and care.
- » For moderate risk:
 - Use the H.O.P.E. actions. This means looking for hints, asking directly about suicide, validating pain, and creating a safety plan, including exploring reasons to live.
 - Recommend a mental health assessment within 24–48 hours.
- » For higher risk:
 - Recommend a mental health assessment immediately.
- » As we reviewed in an earlier module, the C-SSRS is a tool that offers guidance for identifying risk of suicide. Refer to **Handout 4A: Columbia-Suicide Severity Rating Scale—Screen with Triage Points for Victim Advocates**.

Handout 4A: Columbia-Suicide Severity Rating Scale – Screen with Triage Points for Victim Advocates

Ask questions that are bolded and <u>underlined</u> .	Past month	
	YES	NO
Ask questions 1 and 2		
1. <u>Have you wished you were dead or wished you could go to sleep and not wake up?</u>		
2. <u>Have you had any actual thoughts of killing yourself?</u>		
If YES to 2, ask questions 3, 4, 5, and 6. If NO to 2, go directly to question 6.		
3. <u>Have you been thinking about how you might do this?</u> e.g. "I thought about taking an overdose but I never made a specific plan as to when where or how I would actually do it...and I would never go through with it."		
4. <u>Have you had these thoughts and had some intention of acting on them?</u> As opposed to "I have the thoughts but I definitely will not do anything about them."		
5. <u>Have you started to work out or worked out the details of how to kill yourself? Do you intend to carry out this plan?</u>		
6. <u>Have you ever done anything, started to do anything, or prepared to do anything to end your life?</u> Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.		
If YES, ask: Was this within the past three months?		

Possible Response Protocol to C-SSRS Screening

- Item 1 Behavioral Health Referral
- Item 2 Behavioral Health Referral
- Item 3 Behavioral Health Consult (Psychiatric Nurse/Social Worker) and consider Patient Safety Precautions
- Item 4 Behavioral Health Consultation and Patient Safety Precautions
- Item 5 Behavioral Health Consultation and Patient Safety Precautions
- Item 6 Behavioral Health Consult (Psychiatric Nurse/Social Worker) and consider Patient Safety Precautions

Source: Columbia Lighthouse Project, 2020



- » The yellow is considered lower risk, orange is moderate risk, and red is higher risk. As with all of the work we do, it is essential to trust your professional judgment. If a person seems at lower risk according to the screening triage, yet, your professional judgment indicates the risk is higher, you would act upon the higher risk. This is particularly true with crime victims, as there are many factors to consider, including trauma and precipitating factors related to recent victimization.
- » Screening Guidance:
 - Higher risk:
 - Suicidal thoughts with desire to act on it or with a plan
 - Any suicidal behavior in the past 3 months
 - Moderate risk:
 - Suicidal thoughts with no plan, desire to act on it
 - Suicidal behavior more than 3 months ago
 - Multiple risk factors and limited protections against suicide
 - Lower risk:
 - Wish to die, yet strong protections against suicide
 - Risk factors that are changeable
 - No reported history of suicidal thoughts or behavior
- » A mental health assessment is needed to do a thorough interview and determine the appropriate level of care and a treatment plan. No study has identified one specific risk factor as specifically predictive of suicide or suicidal behavior. Therefore, at the end of a suicide assessment, an estimation of suicide risk is based on sound judgment. It is important to keep in mind that this is guidance for responding to screening, and screening is only one part of the process.
- » Once you have screened for and identified risk, recommended actions include referring the person to a mental health professional to assess risk, engaging in treatment, and establishing a safety plan.
- » We've looked for hints, asked openly about suicide, and now we'll focus on **validating pain**. Sometimes, we want to move quickly and solve the problem. Yet, victims need to be heard. They need to know that you really understand their pain.

H.O.P.E. Saves Lives



Look for Hints



Ask Openly about suicide



Validate Pain



Explore reasons to live and plan to stay safe

- » **Reflection:** What are some ways you have validated a victim's pain?
- » It can be hard to know how to respond when a victim communicates that they are thinking about suicide. Often, our reactions, such as fear or insecurity about how to respond, can become our internal focus. Take a deep breath. Validate their pain. Communicate that you have truly understood and heard their pain. There are times this requires silence and pausing. It takes a lot of courage for persons to be vulnerable enough to share that they are experiencing so much pain, that they are considering suicide. Connect with them. Commend them for speaking up and acknowledging their suffering to another person. Validate not only the pain related to their suicidal crisis, but their experience as a victim. After you have validated their pain, then you can offer hope and refer them to engage in treatment. For example, you might say, "Things have just been so overwhelming for you. It seems as if you really want a way out of this intense pain. Thank you for telling me how difficult things have been. I will support you through this time. You don't have to go through this alone. I think there is a way to work through this experience and gain some relief from your pain. Let's think about the part of you that wants to live and get you some support and help to decrease your pain."
- » These are some sample statements that you can say when communicating with a victim.
- » It is important to connect, without making overly simplified statements. These are examples, but as we all know, one line will not resolve a suicidal crisis:
 - "I'm here with you and want you to get through this."
 - "I know things are really hard right now and feel overwhelming."
 - "I think with some help and resources we can get you support through this difficult time."
 - "It sounds like you have been feeling very down and stressed. I want to make sure you get support to get you through that stress."

- » As a part of the development of this training, Education Development Center asked advocates to share what they would say to a victim struggling with suicidal thoughts. Take a couple minutes to watch this video: <https://edc.org/hopeforadvocates>. You can also share this with others in your workplace. Think about what you would add and how you would validate pain for a person at risk for suicide.
- » The person in the suicidal crisis is often ambivalent—part of them wants to live, part of them wants to die. Validating their pain lets them know you have truly heard them.
 - Exploring reasons to live connects with the part of them that wants to live. This can also be a way to strengthen support and the things helping to protect them from suicide.
 - Most persons who think about suicide will not attempt or die by suicide.
 - Gently re-focusing to consider the choice of life can help to equalize emotions, remind them of choice, and open up a conversation about new types of help/support that they may not have tried or considered.
 - Focus on the person’s values and perspective and make linkages to areas of their life you know to be important.
 - Most persons who think about suicide will not attempt or die by suicide. By using these skills, we can help victims experience recovery and healing.
- » **Reflection:** What are some statements you might say to explore reasons to live and offer hope?
- » Let’s hear from a voice of recovery and healing. We’ll hear Shelby Rowe share her pain that led to a suicide attempt. The video will end with her own statements that she would have said to herself to validate pain and provide hope: <https://www.usatoday.com/videos/news/2018/11/20/suicide-survivor-profile-shelby-rowe/1737400002/>.
- » After we explore and engage the part of the person that wants to live, we can begin to collaboratively establish a safety plan. We will be discussing the specifics of a safety plan a little later. Part of a comprehensive safety plan is engaging in a mental health assessment and treatment. This includes knowing your local mental health resources. These are some of the resources that you can use to create a safety net. It is best to have these phone numbers and locations in your back pocket and ready when you need them. **Handout 4B: Resources** has a list of national crisis resources.

Handout 4B: Resources

This is a list of national resources that can be useful to you, as the advocate, and to a person in a suicidal crisis. In addition to these resources, there are numerous tribal and local crisis, emergency, and treatment services.

<p>National Suicide Prevention Lifeline National network of local crisis centers providing 24/7 free, confidential support</p>	<p>Phone: 1-800-273-8255 For deaf/HoH: 1-800-799-4889 via TTY En Español: 1-888-628-9454 Online chat: https://suicidepreventionlifeline.org/chat/ Website: https://suicidepreventionlifeline.org/</p>
<p>Veterans Crisis Line 24/7 free, confidential support for veterans and their loved ones (caller does not have to be registered with the VA or enrolled in VA health care.)</p>	<p>Phone: 1-800-273-8255 [Press 1] For deaf/HoH: 1-800-799-4889 via TTY Text: 838-255 Website & online chat: https://www.veteranscrisisline.net/</p>
<p>Crisis Text Line 24/7 free support from crisis counselors through text messaging</p>	<p>Text: Text HOME to 741-741 Website: https://www.crisistextline.org/</p>
<p>Man Therapy Resources about anger, stress, sadness, substance abuse, and suicide geared toward men</p>	<p>Website: http://mantherapy.org/</p>
<p>Local resources (write in with resources in your area)</p>	<p>Primary care provider _____ Local psychiatric hospital _____ Local walk-in clinic _____ Local emergency department _____ Local urgent care center _____</p>
<p>MY3 Suicide prevention app that allows users to add three people to talk to when in crisis, build a safety plan, and access resources</p>	<p>Apple App Store: https://itunes.apple.com/us/app/my3-suicide-lifeline/id709651264?mt=8 Google Play: https://play.google.com/store/apps/details?id=com.nerdery.my3 Website: http://my3app.org/</p>

<p>Now Matters Now Collection of videos that teach Dialectical Behavior Therapy (DBT) skills to address suicidality and other issues</p>	<p>Website: https://www.nowmattersnow.org/skills</p>
<p>Safety Planning Intervention Information about creating safety plans for those dealing with suicidal thoughts</p>	<p>Website: http://www.suicidesafetyplan.com/</p>
<p>The Trevor Project Free crisis line, chat, and text for LGBTQ youth (ages 13–24)</p>	<p>Phone: 1-866-488-7386 [24/7] Text: TREVOR to 1-202-304-1200 Website and online chat: https://www.thetrevorproject.org/</p>
<p>Trans Lifeline Hotline by and for transgender people</p>	<p>Phone: 877-565-8860 Website: https://www.translifeline.org/</p>
<p>RAINN National Sexual Assault Hotline 24/7 free, confidential support for survivors of sexual assault</p>	<p>Phone: 800-656-HOPE (4673) Find a local service provider: https://centers.rainn.org/ Website and online chat: https://rainn.org/</p>
<p>Love Is Respect 24/7 support for teens involved with dating abuse</p>	<p>Phone: 1-866-331-9474 Text: LOVEIS to 22522 Website and online chat: http://www.loveisrespect.org/</p>
<p>Victim Connect National hotline for crime victims</p>	<p>Phone: 1-855-4VICTIM (1-855-484-2846) Mon.–Fri.: 8:30 a.m.–7:30 p.m. Website: https://victimconnect.org/</p>
<p>National Domestic Violence Hotline 24/7 hotline for people experiencing domestic violence</p>	<p>Phone: 1-800-799-7233 For deaf/HoH: 1-800-787-3224 via TTY Website: http://www.thehotline.org/</p>



- » When responding to risk, immediate referral and resources can include:
 - Community mental health
 - Tribal health and mental health services
 - Mobile crisis
 - Existing treatment provider(s)
 - Emergency department
 - Inpatient assessment
 - Suicide prevention crisis lines
 - Local
 - National
 - Specific populations: Veteran/military, LGBTQ
 - Other crisis supports, such as RAINN and domestic violence crisis support
- » How do you know what the best referral is? It depends on what the problem is, the needs of the individual, their situation, and supports available in their community. For example, is there complex trauma, substance use issues, or immediate suicide concerns? Knowing your resources and considering general guidance, use your judgment and consultation for what is best for the person. You can do this in collaboration with the victim and with their support system. Collaboration is a way of empowering the victim and helping them to recover a sense of control. Healing can be facilitated through shared power and decision-making.
- » We know that many of you may be in situations of responding to a person at risk in a variety of settings. For example, you can use screening and engagement with a victim over the phone. If you are talking with someone at risk via text or email, it is important to consider the best way of continuing to engage them. This could be via electronic communication for some people. For others, it could be better to make a call or connect with them for an in-person meeting.
- » It is important to engage support and communicate any safety and risk concerns to anyone who may be providing treatment or assisting. Throughout this manual apply these recommendations to your specific settings and communities.
- » If the person is already engaged in treatment, communicate with existing treatment providers the necessary information regarding the person's safety and suicide risk. This can be done through release of information, collaboratively engaging the person at risk to share information. It is essential to share only the information that the mental health treatment provider needs and to maintain privacy and confidentiality.
- » If a person says no to engaging in care, especially if they have strong protective factors, engage support and strengthen protective factors. It is okay to trust your gut, especially if cultural norms discourage talking about a suicidal crisis. Even if a victim has things that

protect them from suicide, it is still important to connect them to care. Remember, you don't have to do this work alone. You can seek consultation and run it past a colleague or mental health professional and engage the person's support system.

- » Almost all efforts to help someone to live, reduce their pain, and not attempt suicide will usually be met with agreement and relief. Don't hesitate to get involved or take the lead. For some of you, it may be your role to **establish a safety plan** specific to suicide with the victim. For others, you might support the safety plan that a mental health professional creates.

H.O.P.E. Saves Lives



Look for Hints



Ask Openly about suicide



Validate Pain



Explore reasons to live and plan to stay safe

- » A safety plan is a prioritized written list of strategies and resources that victims can use in times of distress (Stanley & Brown, 2012).
- » A safety plan is brief, in the person's own words, and easy to read. The title of a safety plan can be adjusted to fit the community and individual. With some individuals or communities, it may be best to call this a healing plan or some other title that fits for the person at risk for suicide. For this discussion, we are focused on a safety planning intervention that was developed by Barbara Stanley and Greg Brown. As we go through this, think about how you might adjust the language or add to this plan to make it specific to your community and the individuals you serve.
- » The process of safety planning is done through validating and engaging the person. Collaborative engagement is key to the likelihood that the safety plan will be used. There are six steps involved in the development of a safety plan.
- » An example safety plan for suicide risk is included in **Handout 4C: Patient Safety Plan Template**.

Access the safety planning form and training manual for the suicide-specific safety planning intervention at Safety Planning Intervention: A brief intervention for reducing suicide risk: http://suicidesafetyplan.com/Page_8.html.

A similar evidence-based tool is Crisis Response Planning for Suicide Prevention, which can be accessed at <https://crpforsuicide.com/>.

Handout 4C: Safety Plan Template

The safety plan can also be called a wellness plan or another title which would connect more with an individual at risk for suicide and their support system.

Step 1: Warning signs	
1. _____	
2. _____	
3. _____	
Step 2: Internal coping strategies – Things I can do to take my mind off my problems without contacting another person	
1. _____	
2. _____	
3. _____	
Step 3: People and social settings that provide distraction	
1. Name _____	Phone _____
2. Name _____	Phone _____
3. Place _____	
4. Place _____	
Step 4: People whom I can ask for help	
1. Name _____	Phone _____
2. Name _____	Phone _____
3. Name _____	Phone _____
Step 5: Professionals or agencies I can contact during a crisis	
1. Mental Health Professional name Mental Health Professional emergency contact #	Phone _____
2. Mental Health Professional name Mental Health Professional emergency contact #	Phone _____
3. Suicide Prevention Lifeline: 1-800-273-TALK (8255)	
4. Local emergency service _____ Emergency services address _____ Emergency services phone _____	
Making the environment safe	
1. _____	
2. _____	
<small>Safety Plan Template, ©2008 Barbara Stanley and Gregory K. Brown, is reprinted with the express permission of the authors. No portion of the Safety Plan Template may be reproduced without their express permission.</small>	

Source: Stanley & Brown, 2008

- » For Step 1, which is focused on warning signs, ask, “How will you know when the safety plan should be used?” “What do you experience when you start to think about suicide or feel extremely depressed?” Then list warning signs and triggers, which can include thoughts, images, moods, and/or behaviors, using the person’s own words (Stanley et al., 2009).
- » In Step 2, the person outlines internal coping strategies. Ask, “If you are on your own and you become suicidal again, what can you do to help yourself to not act on your thoughts or urges?” This might include traditional and spiritual practices (Stanley et al., 2009).
- » Identify the likelihood that the person will use the safety plan by asking, “How likely do you think you would be to do this step during a time of crisis?” Ask about potential obstacles by asking, “What might stand in the way of you thinking of these activities or doing them if you think of them?” Use a collaborative, problem-solving approach to address potential barriers and identify alternative coping strategies (Stanley et al., 2009).
- » Step 3 can be used if the coping strategies do not resolve the suicidal crisis. For this step, ask, “Who or what social settings help you take your mind off your problems, at least for a little while?” and “Who helps you feel better when you socialize with them?” Identify safe places where they can go to be around people (e.g., a coffee shop) at a time of crisis. It is key that these are accessible when the crisis typically happens. So, walking at around a store likely could not be used for a crisis at 2:00 a.m. Engage the person in listing several people and social settings to increase their options. The goal is distraction from suicidal thoughts and feelings. As with the other steps, consider the likelihood the victim will use these supports, identify possible barriers, and problem solve as needed (Stanley et al., 2009).
- » In Step 4, the person identifies people who could offer support at a time of crisis. Ask, “Among your family or friends, who do you think you could contact for help during a crisis?” or “Who is supportive of you, and who do you feel that you can talk with when you’re under stress?” This might include a person that the victim sees as a leader in their nation or community. Someone they look up to and feel safe to talk with. As with Step 3, identifying more than one person is helpful. Aim to list at least three support people, then prioritize the list. In this step, a person would share that they are in crisis to others. Again, consider the likelihood the victim will engage these supports, identify possible barriers, and problem solve. Use of role-play and rehearsal can be very useful as well (Stanley et al., 2009).
- » In Step 5, the victim identifies professional supports. You can also add and educate the victim about professional and crisis supports. Ask, “Who are the mental health professionals and crisis supports that we should identify to be on your safety plan?” This would include traditional healers and medicine. List specific information, such as names, numbers, and/or locations of professionals or local urgent care services (Stanley et al., 2009).
- » Step 6 is about making the victim’s environment safe, especially if they are at a higher risk of suicide. Limiting access to lethal means, for example, can help keep a person

safe in a time of intense crisis. It is a way to add some time and space for the person who might have progressed on the pathway to suicide. Reduced access to lethal objects, such as safely storing a firearm, can help to keep a person safe in a time of intense crisis. However, there are unique considerations with a crime victim (Stanley et al., 2009).

For example:

- In Native communities, we know that if a person engages in suicidal behaviors, they might be more likely to use a method such as hanging. While this may be harder to reduce access to, it can be important to reduce access to someone's preferred means. This might include a preferred belt, favorite rope, or preferred location.
 - Some victims may be concerned about their physical safety and so choose to own a firearm to protect themselves from others. This very real safety concern must be balanced with the victim's safety in a time of crisis. In some situations, it may be appropriate to have the firearm in a locked safe, use a trigger lock, or store the ammunition and firearm in different locations.
 - In many other situations, something like a photo or something that is representative of a protective factor, such as traditional and healing images and objects, children, spiritual imagery, pets, or others, can help protect a person in a suicidal crisis.
- » Ask which means the person would consider using during a suicidal crisis: "What things do you have access to and may use to attempt to kill yourself?" Then with the person, collaboratively identify ways to secure or limit their access to lethal means. Ask, "How can we go about developing a plan to limit your access to these?" This could include storing medications in a lock box, keeping only a small supply in the home, or removing an item the person identifies. Research has shown this practice to be effective, and that often, persons don't substitute means. So, if the identified or desired method is limited or secured, the person may be safer in a time of crisis (Stanley et al., 2009).

For clients who are experiencing domestic violence, it is crucial to interweave support and resources for domestic violence with the suicide support. Both must be addressed. In addition, for clients who are experiencing domestic violence and not planning to leave their partner, offering resources for counseling/intervention for the person who is using violence against them is an important and, often, overlooked step. When these clients are referred to a mental health provider, it is important that the provider is knowledgeable of both suicide AND domestic violence.

- » We know that our communities and individuals are resilient. We know that we've been resilient for generations. It's in our genes and our lands.
- » There is hope. And suicide deaths can be prevented. Research has shown that simply receiving a postcard from staff for individuals who were treated in an inpatient hospital for a suicide attempt actually decreased suicide deaths (Stanford SPARQ, n.d.; Zero Suicide Institute, 2020a).

- » Follow up in a way that is supportive and collaborative:
 - Reach out and continue to show support.
 - This can be through a text, a phone call, an email, or a letter.
 - Caring contacts, ongoing supportive messages that don't ask for anything or remind about an appointment, have been shown to save lives (Doupnik et al., 2020).

Safer Suicide Triage

Introduction

This section focuses on the process when a victim advocate has identified a client who may be at increased risk of suicide. Some settings will have savvy clinicians on-site who can provide treatment for suicide risk, but most sites will not have these resources.

Now that we have addressed how to respond to someone who may be at risk for suicide, it is time to discuss how to figure out the next step to ensure they receive care.

Objectives

Participants will:

- » Increase awareness of the types of practice-based interventions available for suicide to be able to describe options to victims
- » Develop a framework for transitioning a person at risk for suicide to the next level of care

Key Points

- » Caring for people at risk can mean a lot of things. For our training, it means supporting as an advocate, caring within the tribal ways, traditional medicine, and using best practices.
- » Evidence-based suicide-specific treatment:
 - Mental health care that directly targets and treats suicidal thinking and behaviors along with the contributing and protective factors to suicide risk is essential for effective treatment. This treatment addresses the suicide risk and behavioral health disorders alongside care for trauma using best practice and promising treatments. For a person in a suicidal crisis, it is not sufficient only to address behavioral health concerns such as depression, anxiety, or post-traumatic stress.

- Best and promising practices using suicide-specific treatments include:
 - Dialectical Behavior Therapy (DBT)
 - Cognitive Behavioral Therapy for Suicide Prevention (CBT-SP)
 - Collaborative Assessment and Management of Suicidality (CAMS)
 - Non-demand follow-up contacts, such as letters and phone calls (Zero Suicide Institute, 2020b)

For additional resources regarding safer care for persons at risk for suicide reference, see the toolkit *Best and Promising Practices for the Implementation of Zero Suicide in Indian Country* at <http://zerosuicide.edc.org/toolkit/indian-country>.

- » Additional examples of culturally relevant interventions that can be partnered with those that are evidence-based include:
 - [Honoring Native Life: Creating Conversation Around Suicide Prevention & Response](#)
 - [Gathering of Native Americans \(GONA\) Training of Facilitators](#)
 - [A Journey Home: A Canoe Movement in Southeast Alaska](#)
 - [WeRNative: For Native Youth, by Native Youth](#)
 - [#ThereIsHope—Suicide Prevention and Awareness on the Pine Ridge](#)
 - [Project Venture: Adventure With an Indigenous Mind](#)
 - [The Sky Center: Natural Helpers, A Peer-Helping Program](#)
 - [Suicide Prevention: A Culture-Based Approach in Indian Country](#)
 - [Hope on the Reservation: Combating Suicide](#)
- » As an advocate, your role is to be knowledgeable of who provides these evidence-based treatments in a trauma-sensitive context and make referrals, as necessary. In some regions, options might be limited, and it may be important to understand who specializes in treatment of suicide risk and trauma. As with all of our work, referrals to and the provision of treatment should be culturally appropriate and person-centered.
- » Considering how you would triage when responding to a crisis or identifying a victim at risk for suicide, complete **Worksheet 4D: Mental Health Supports** with resources available in your region specific to emergency contacts and crisis contacts, routine mental health contacts, and persons to go to for consultation:
 - In some areas, the mental health contacts may all be in the same agency, or there may be only a few agencies. In more urban or resourced areas, there could be 24-hour treatment facilities, mobile crisis services, inpatient hospitals, and a variety of routine mental health facilities.
 - As you complete the contacts and processes for your organization and role, if there is something that you cannot complete right now, make a note of the person you will need to contact to find the answer to complete the process.

Worksheet 4D: Mental Health Supports

Emergency and Crisis Mental Health Contact(s)

Urgent Mental Health Contact(s)

Routine Mental Health Contact(s)

Person(s) to go for Consultation




Conclusion

Key Points

- » This section helped to bring all of the sections together. When responding, we need to be aware of our reactions and the impact they may be having on the effectiveness of our responses and how in sync we are with the person at risk. In order to respond effectively, we need to bring together what we know about empowering our communities, the suicidal crisis, and the intersection with victimization. Now, we have the resources and ability to respond in a way that is collaborative, genuine, and supportive.
- » We have covered a lot of content in the training at this point. In the next section, we will discuss caring for ourselves. This component is essential to support our work. It is similar to putting a plane's oxygen mask on yourself first before assisting others. Caring for ourselves and our colleagues allows us to be in the zone, prevents burnout, and enables us to support our teammates.

Module 5: Recharge Ourselves



Look for Hints



Ask Openly about
suicide



Validate Pain



Explore reasons to live
and plan to stay safe

H.O.P.E. saves lives

Module 5 Overview

+ Topics Covered

- » Personal and professional self-care
- » Models of self-care
- » Processing a suicide loss



+ Learning Objectives

In Module 5, participants will:

- » Recognize one's professional obligation to attend to self-care
- » Identify the warning signs of burnout and vicarious trauma
- » Describe at least three skills that are particularly helpful in supporting self-care
- » Identify resources for coping with a suicide loss



+ Materials

- » **Worksheet 5A: Self-Care Reflections**
- » **Worksheet 5B: My Self-Care Plan**



Introduction

Introduction

The work we do day to day is very different from what most people do for a living. Many of us came to this field because of our passions, experiences, and love for our communities. We put our all into supporting those impacted by crime, working to keep people and communities safe, and providing the best opportunities for recovery, healing, and thriving. Often, this means long hours and hearing difficult stories. This dedication and focus can impact our personal lives. It is demanding work, and at times, it can feel as if it is thankless work. But our work—your work—matters, and it makes a difference. Self-care and community care are absolutely necessary for your safety, your health, the health of your relationships, the strength of your community, and the effectiveness of your work.

Commitment to Self-Care

Introduction

Often, the topic of self-care is discussed solely as something that is “good to do” for oneself. This is not the whole picture. We know that professionals who do not take care of themselves and who do not get enough rejuvenation to restore their mental flexibility can actually harm their clients. Of course, no one in this field intends to harm their clients. Yet, when we are not healthy:

- » We make poor decisions about maintaining boundaries.
- » We ask fewer questions and do less thorough assessments.
- » We can appear less empathic and caring to our clients. This can result in them losing hope that there is anyone who can help them.

So, the first thing to realize about self-care is that it is a professional responsibility—not simply a good thing to do for yourself.

It is important to recognize that caring for and recharging yourself is essential to your role. It is your responsibility to your work, your nation, and your community you are supporting.

Objectives

- » Recognize one's professional obligation to attend to self-care

Reflection

- » Complete **Worksheet 5A: Self-Care Reflections**

Worksheet 5A: Self-Care Reflections

Reflection Questions

1. Think back to a time when you felt like you were really in sync with the people you work with and felt like things were going really well. What was your self-care and/or support like at the time?
2. Think back to a time when you crossed an ethical or a professional boundary or came close to doing so. Examples can include when you became attached to a client in some way that felt too strong and/or unhealthy for you. What was your self-care and/or state of mind like at the time?
3. How does caring for yourself impact your work and the people you work with?
4. What happens to the people you work with and the community you support if you don't take care of yourself?
5. Why is it important for you to recharge and care for yourself?



Vicarious Traumatization

Introduction

Let's review some definitions and common signs of burnout and vicarious trauma. Then, we will shift to ways to care for yourself. The signs that we review are not an exhaustive list. You know yourself best. You know your team and colleagues best. Use this expertise to strengthen yourself and your team.

Objectives


- » Identify the warning signs of burnout and vicarious trauma
- » Describe at least three skills that are particularly helpful in supporting self-care
- » Identify resources for coping with a suicide loss

Key Points

- » Burnout is a job-related stress condition. It typically progresses over time and is often a result of chronic stress or frustration. In the fields that we are working in, burnout can occur in a shorter period due to the intensity of our work.
- » Signs of burnout can be specific to an individual and can be impacted by culture. Some signs include feeling emotionally exhausted, separating oneself, and having a reduced sense of accomplishment (Institute for Quality and Efficiency in Health Care, 2017). Typically, the drive that pushes us to go above and beyond is impacted by burnout. We can find ourselves less passionate about our work and detached.
- » Vicarious trauma is a state of tension and preoccupation with the stories/trauma experiences described by Clients (American Counseling Association, n.d.). We work closely with trauma survivors and can empathically take on aspects of the trauma that the survivor experienced. This can result in experiencing symptoms similar to those of the trauma survivors themselves. The signs of vicarious trauma can look like post-traumatic stress disorder symptoms.

It's important to stress that vicarious traumatization is a normal response to doing the hard work of supporting and advocating for people who have experienced trauma and violence and is not a reflection of the advocate's competence. Just as there are individual differences in how people respond to traumatic life events, there are also individual differences in what leads to vicarious trauma in the helper (Schauben & Frazier 1995).

- » The *Vicarious Trauma Toolkit* (VTT) was developed based on the idea that exposure to the traumatic experiences of other people is an inevitable part of the work that occurs in the field of victim services (Office for Victims of Crime [OVC], n.d.b). However, organizations can reduce the impact of the negative effects of trauma exposure by becoming informed about vicarious trauma. The VTT includes tools and resources that were created for organizations to support the needs of their staff (OVC, n.d.a).



Source: Office for Victims of Crime, n.d.c

- » While we can all take individual responsibility for our self-care, organizations and communities have a responsibility in caring for staff (OVC, n.d.a).
- » Take a moment to consider: What do you think are some organizational factors that can help prevent burnout and vicarious trauma among staff?
- » Several organizational factors that might prevent burnout and vicarious trauma include:
 - Having a safe environment to talk to colleagues and ask for help or support.
 - Routinely building in support and helping each other.
 - De-briefing in a timely and supportive manner.
 - Being able to take time off when needed—not just when seeing the signs of burnout, vicarious trauma, or toxic stress, but also being able to take time off for vacations, to recharge, and to take care of personal matters.
 - Being able to access professional support when needed, including Employee Assistance Programs.

- Having a balance in working hours—knowing that sometimes, work will be quite intense and may require overtime, while at other times, you may have less demands. A balance in workload is important, especially if you have been working with several particularly challenging cases.
 - For instance, if you have been working with many crime victims who have experienced intense physical violence, and you have started to feel overwhelmed by the severity of the cases, you might take on a different case, which, while still challenging, is not specific to physical violence.
- » Let's talk about individual ways to support your self-care. There are many self-care strategies. Common categories for self-care include:
 - Physical health promotion activities
 - Spiritually oriented activities
 - Leisure activities
 - Seeking emotional support
 - Decompression strategies/rituals
 - Setting boundaries around work
 - Setting boundaries about types of information ascertained from clients
 - Others (write in your ideas): _____

Resources

Introduction

At times, in this work, we might lose someone to suicide. It is important to take stock of the effect that a suicide loss can have on your own sense of well-being, as well as on your colleagues, your organization, and on the family and larger community. We will focus the next section on taking care of yourself and on strategies to implement for postvention care of a team. We will discuss ways to support family members in Module 7.

Key Points

- » If you are impacted by a suicide loss, remember everyone copes and responds in their own way. There is no right way to grieve or approach a loss that might impact your professional work. It may be helpful to do the following:
 - Take time to become aware of your needs and continue to do so.

- Seek support:
 - From colleagues & support system
 - Support groups
 - Professional therapy
 - Blogs, listservs, online support
- Ask for help, particularly if you are assigned a similar case.
- Take the time you need.
- Consult.
- » If grief (or another emotion) is affecting anyone's work, personal relationships, or ability to sleep, eat, or engage in daily activities, likely additional support may be needed.
 - Seek professional treatment.
 - Add additional peer, family, spiritual, or other help.
 - Leverage consultation and team support as you are able.
 - Consider taking a break, such as stepping back from your role for a little while or taking a vacation.
 - Use Family Medical Leave Act (FMLA) time for treatment and additional support.
- » Remember to honor all emotions. Anger at yourself for being unable to prevent the loss and/or anger at the client for taking their life are normal emotions, and they require time and space to work through.

Recharge Ourselves

Objectives

Participants will:

- » Describe at least three coping skills that are particularly helpful in supporting self-care

Key Points

- » Identify strategies for self-care.
- » Focus on applying ideas to day-to-day life.

Reflection and Activity

- » Let's shift to identifying and practicing how we can care for ourselves. While completing **Worksheet 5B: My Self-Care Plan**, think outside of the norm. For instance, exercising or socializing can be great strategies. Also, think about the times you struggle the most during the day. How might you use strategies during these times? Think about how you might be able to use the drive to and from work to practice strategies or how you might practice calming yourself just before you go to sleep.
- » First, write personal signals of burnout or vicarious trauma.
- » Then, identify personal strategies, social supports, and professional support you can use.

Worksheet 5B: My Self-Care Plan

What signals to me that I am beginning to struggle and that I need to focus on my own wellness? These include thoughts, feelings, physical sensations, and actions.

What strategies can I use to positively impact my wellness?

Who can I contact among my community and nation, including my family and friends, to help me?

What traditional healing practices can I use to support my health?

If I would like to seek professional help, what professionals are available to me?




- » Now that we have identified our personal signals and self-care strategies, I hope you can commit to implementing at least one of these today and several in the next week.
- » You can take a few minutes for self-care right now.
- » If you would like, you can engage in this guided breathing exercise.
 - Setting your guide, phone, and anything else aside, find a comfortable and upright posture that supports your body and breathing. Place both feet on the floor, arms uncrossed, if you are able to do so. You might use the back of the chair for support. You can keep your eyes open or closed for this exercise. If you keep them open, look ahead of you just a few feet.
 - Start by just observing your breath wherever you feel it most predominantly. This can be in your chest, lungs, belly, or at the tip of your nose as the air moves in and out.
 - You are not trying to change the breath right now. You're just taking a few moments to become aware of your breath. Put thoughts of the past or plans for what is next off to the side. Just be here, right now.
 - Now, take a few deep inhaleds and exhaleds. Start by inhaling for four counts. 1-2-3-4. Make sure you fill your lungs completely to the point where you feel your abdomen is filled full of air like a balloon. Hold your breath for four counts then slowly exhale for eight counts.
 - It will take a few attempts before you feel comfortable, but you will soon find a rhythm that suits you. There is no wrong way of doing this. Inhale for four counts, 1-2-3-4; hold for four counts, 1-2-3-4; and then slowly exhale for at least eight counts, 1-2-3-4-5-6-7-8. Continue doing this at your own pace.
 - Repeat this exercise several times.
 - Now, return to your normal breathing without trying to manipulate or change it.
 - Just check in with your body, starting with your head. Scan from your head down to your feet. Notice any place you observe tightness or tension. See if you can invite that part of your body to relax. Any time you notice yourself thinking about the past or planning for the future, just return to your body and focus your attention on this.
 - As you scanned your body, you may have noticed areas that need some attention. You may stand, move, or stretch in any way that you personally need it. This could be rolling your shoulders or holding your arms up and stretching to the sky. You can take a few minutes for this.

Conclusion

Key Points

- » Prioritizing one's own well-being is a requirement of your job.
- » The people and communities we serve struggle with significant concerns, and they need supportive advocates who are healthy to help support them in the ways they deserve.
- » Communities and organizations have a responsibility in creating a culture of support and protections against vicarious trauma and burnout. We must think beyond self-care and prioritize the well-being of all staff. This includes building in traditional and cultural healing practices in addition to strategies to support staff.

Module 6: Consider the Complexities



Look for Hints



Ask Openly about
suicide



Validate Pain



Explore reasons to live
and plan to stay safe

H.O.P.E. saves lives

Module 6 Overview

+ Topics Covered

- » Explore the linkage between suicide and trauma, suicide and substance use, and substance use and trauma
- » Gain an understanding of these linkages through real-life scenarios
- » Implications of understanding these linkages, including distinguishing between some substance use behaviors and suicidality
- » Recognize the complexity involved in distinguishing some substance use behaviors from thoughts and feelings suicidal
- » Identify strategies to resolve unclear situations

+ Learning Objectives

In Module 6, participants will:

- » Identify the linkages between suicide, trauma, and/or substance use
- » Recognize the complexity involved in distinguishing some substance use behaviors from suicidal thoughts and feelings
- » Identify strategies to resolve unclear situations

+ Materials

Worksheet 6A: [Reflection – Examples](#)

Worksheet 6B: [Unclear Situations](#)

Video: [Johnathan White-Cloud Courtney Shares His Experience](#), (start at 3:20; end at 6:18; 2:58 minutes in length): <https://www.nowmattersnow.org/skill/suicidal-thoughts>

Introduction

Introduction

Too often, we in the crime victim arena talk in silos. We have gotten much better at drawing connections between someone's trauma history and their risk for victimization, and in understanding that violent victimization can lead to mental health issues. However, we are often not discussing the linkages between crime victimization, trauma, suicide, and/or substance use. If we are not aware of the interconnections, we will miss hints and opportunities to help people address their underlying pain. Far too often, we become sidetracked by sending someone to substance use treatment or thinking someone doesn't want to change because they continue to abuse substances when, in reality, their underlying emotional pain is so great that substance use is the only way they know to cope. Alternatively, sometimes we send someone to the emergency department (ED) because we are afraid of suicidality and do not take the time to understand how the experience of trauma, including generational trauma and/or substance use, is related to their current suicidal crisis. As such, those pieces of the puzzle never get documented. Thus, once the ED staff believe the person is stable from a suicide perspective, they are released, and there is no follow up to the underlying issues. This module focuses on the interconnections between trauma, suicide, and substance use.

Key Points

- » Identify cyclical and reinforcing linkages between suicide and trauma and/or substance use.
- » Recognize the complexity involved in distinguishing some substance use behaviors from suicidality.
- » Identify strategies to resolve unclear situations.

Consider the Complexities

Objectives

Participants will:

- » Identify the linkages between suicide, trauma, and/or substance use
- » Recognize the complexity involved in distinguishing some substance use behaviors from suicidal thoughts and feelings

Key Points

- » The American Psychological Association defines *trauma* as “an emotional response to a terrible event like an accident, a rape, or a natural disaster” (American Psychological Association [APA], 2020). Responses to trauma may include shock, denial, flashbacks, physical symptoms, and unpredictable emotional reactions (APA, 2020).
- » Two people may experience the same event differently—one may experience it as a trauma, and another may not. There is a subjective element to defining a traumatic event.
- » **Reflection:** Think about the difference between stress and trauma. What do you see as the differences between stress and trauma?
- » Everyone experiences stress. Not everyone experiences a trauma.
- » Even among those that experience an event, a set of events, as harmful or life threatening, many people don't have lasting adverse effects, and many people heal on their own.
- » Some types of traumas are associated with a higher likelihood of mental health disorder and/or substance use—usually the more personal and the more severe the type of violent experience, the more significant the impact and greater potential for debilitating consequences.

Individual trauma:

- » Individual trauma includes common traumatic events, such as witnessing a death; being threatened with death, injury, or sexual violence; or experiencing a natural disaster. Only a small percentage of people who experience a trauma develop unhealthy coping habits or have symptoms similar to post traumatic stress disorder (PTSD).

Historical and intergenerational trauma:

- » Historical or intergenerational trauma occurs as a result of decades and centuries of discrimination, violence, and victimization. It happens to populations, often paired with other adversities. Generations of Native people have experienced decades of trauma that have, over time, impacted health and family dynamics for generations.

Suicide and trauma:

- » Adverse childhood experiences (ACEs) include stressful or traumatic events that can lead to social, emotional, and cognitive impairment, which, in turn, can lead to high-risk behaviors, disease, and early death. Felitti and his colleagues first identified the term ACEs when looking at risk factors for chronic disease in a sample of hospital patients (Felitti et al., 1998). Since their initial work, interest has grown in understanding which adverse experiences have what types of long-term health effects. Individuals who reported four or more ACEs had a 12-fold increased risk for attempting suicide (Ports et al., 2017).

- » ACEs scores may not fully reflect additional traumatic experiences such as racism and discrimination. So, as we consider the experiences that are measured with an ACEs score, we also need to keep in mind that there could be other complicating factors.
- » Individuals diagnosed with PTSD have been found to be at higher risk for suicide attempts even after controlling for physical illness and mental health disorders (U.S. Department of Veterans Affairs, 2019). Some studies indicate PTSD is a precipitating factor of suicide. This data emphasizes the intersection of suicide risk and trauma. Victimization that results in physical, sexual, and emotional impacts and develops into PTSD symptoms can lead to a higher risk for suicide.
- » People who do experience early adversity are not fated to experience problems. We know from both experience and research that with the right kinds of support, most people can heal.

Suicide and substance use:

- » Research indicates there are clear linkages of trauma leading to substance use, substance use leading to trauma, and the circular nature of both existing together. Substance use disorders may come before trauma exposure. ACEs have a dose-response relationship with many health problems. As researchers followed participants over time, they discovered that a person's cumulative ACEs score had a strong relationship to numerous health, social, and behavioral problems throughout their life span, including substance use disorders (SAMHSA, 2018).
- » Preventing suicide death will also require an understanding of the complexities with substance misuse (Oquendo & Volkow, 2018). At times substance overdose deaths are ruled intentional and may actually have had suicidal underpinnings. It can often be very difficult, at times impossible, to determine the difference between an unintentional drug overdose and a suicide attempt. Yet, making a distinction can have impacts on treatment decisions. Often, staff may be quick to label "overdoses" and thus may inadvertently ignore suicidal behavior or ambivalence about living.
- » Substances affect thinking and emotional processes and can contribute to a suicidal crisis.
- » In 2011, approximately 230,000 ED visits resulted from drug-related suicide attempts, with almost all involving a prescription drug or an over-the-counter medication. One of the reasons that alcohol and/or drug misuse significantly affects suicide rates is the reduced inhibitions that occur when a person is intoxicated (Pompili et al., 2010). Alcohol and opioids can be a dangerous combination. Currently, less is known about the relationship between suicide risk and other drug use. We know that the number of substances used seems to be more predictive of suicide than the types of substances used (U.S. Office of the Surgeon General & National Action Alliance for Suicide Prevention, 2012).
- » Also, we know that substances are involved in suicide deaths. Of those who died by suicide, 28.3% had problematic substance use; 17% had problematic substance use with alcohol; and 16% had problematic substance use with other substances. It has been

documented that alcohol was present in about 30%–40% of suicide attempts (SAMHSA, 2015).

- » A recent study found that persons using opioids regularly were at greatest risk. They were about 75% more likely to make suicide plans and twice as likely to attempt suicide as persons who did not report any opioid use (Oquendo & Volkow, 2018).

Children Witnessing Domestic and Suicide Risk

A child witnessing domestic violence is linked to suicide risk. Children exposed to chronic domestic violence have a significantly higher lifetime prevalence of suicide attempts. When violence is chronic in a home, even when the child is not directly abused themselves, there is a risk for long-term negative outcomes (University of Toronto, 2016). It is important to remember this when considering the risk for children and early intervention for suicide prevention.

- » The ways that exposure to violence as a child affects our brain development is important to remember when working with victims and understanding the neurological factors that may be contributing to their mental state and even to their suicide risk.

For more information regarding the impact of witnessing domestic violence on a child's developing brain, we recommend watching the video *First Impressions: Exposure to Violence and a Child's Developing Brain*: <https://www.youtube.com/watch?v=brVOYtNMmKk>.

Case Examples and Reflection

- » Trauma can manifest in different ways. Depending upon the trauma experienced and individual and cultural factors, individuals may be drawn to different substances, and the hopelessness related to suicide may play out in a variety of ways. We are going to discuss several examples. As you read these, think about the linkages between trauma, substance use, and suicide risk.
- » **Example:** If someone experiences abuse from a close family member, they may have a hard time attaching to others and creating new, close relationships. They may be overly alert and on guard at all times. Social situations may feel very intense. In addition to the family abuse, they may feel alienated from other Native individuals and their nation. The person may feel isolated trying to acculturate into a culture that is more prevalent around them and separate from their family culture. That person may move toward a substance that allows them to relax and experience less anxiety in social situations. These substances, which can worsen depressive symptoms, may facilitate suicidal behavior.
- » **Reflection:** How can these substances facilitate suicidal behavior? Consider the interconnected and cyclical nature of how trauma history, substance use, and suicide can occur together.
- » If a person is thinking of attempting suicide, alcohol may supply the motivation to act on suicidal thoughts or to complete the steps to die. The victim may believe that alcohol will assist in making dying by suicide painless

- » This becomes a cycle at times, with suicidal behavior or thinking contributing to additional stress and symptoms for the person and serving as a driver to continued substance use.
- » Consider people you have worked with who have struggled with all three of these issues. Think of these examples in relation to the linkages and how suicide, trauma, and substance use are intertwined.
- » We have been discussing complexities focused on substance misuse, trauma, and suicide. There are many factors which can add complexities with the communities and individuals we serve. This includes Native individuals who have served or are serving in the military, young adults who might move from a reservation to urban areas or college, the deep isolation that can exist in some rural areas, and/or the complexities related to technology and the use of social media.
- » In addition to this example and the people you have worked with, let's listen to one person share their experience related to trauma from military experience and a suicidal crisis. This is Johnathan White-Cloud Courtney. He is a member of the Confederated Tribes of Warm Springs, a military veteran, and a suicide prevention and peer support advocate. He has survived suicide attempts, PTSD, and alcohol problems. We will hear him describe his experience. He will talk about hints, warning signs, and triggers to his suicidal crisis and will share what he would say to someone struggling. Watch the video: *Johnathan White-Cloud Courtney Shares His Experience* (start at 3:20; end at 6:18; 2:58 minutes in length) at <https://www.nowmattersnow.org/skill/suicidal-thoughts>.
- » Following the video, pause and consider how you might work with someone like Johnathan to help them towards recovery in all areas. Using **Worksheet 6A: Reflection – Examples**, write down some ways you would work with Johnathan and the other example we reviewed.

Historically, the fields of trauma, suicide, and substance misuse have been separate.

Often, if you seek treatment for substance misuse and are struggling with suicidal thoughts, you might be asked to be stabilized with the suicidal crisis first before being able to get focused substance misuse treatment, and likewise, you might be asked to get clean first before you receive treatment for your trauma. Many professionals trained in trauma and mental health have no substance use training and vice versa. Given this reality, it can be easy for trauma survivors to feel hopeless, increasing the risk present in their lives. Crime victim advocates should be skilled at finding coordinated or integrated care, or helping the client engage with specific service systems as a part of the client's safety plan.

- » The best practice is a combined, holistic, integrated approach.
- » These topics are important to bring together because it is clearly established that people with unresolved trauma histories are much more likely to abuse substances, and that among those with the harder-to-manage substance use disorders, the vast majority have trauma histories.
- » Also, it is clear that those with suicide susceptibility are much more likely to have trauma histories.
- » Our support and help-giving approaches need to align with the evidence and need to address the trauma, substance use, and/or suicide together.
- » If the underlying trauma and substance use aren't considered or addressed, we could invest resources in sending people again and again to ineffective treatment. Meanwhile, they are losing hope in themselves and blaming themselves for not being able to stay clean, when we are giving them the wrong treatment. It isn't fair to ask someone to give up their coping strategy if we haven't helped the huge problem that they are trying to cope with.

Considering a holistic integrated approach, revisit **Worksheet 6A: Reflection – Examples** and write any additional considerations in how you might work with, what you might want to know, and things you would want to consider in the examples reviewed.

Unclear Situations

Introduction

Some situations may be particularly challenging and unclear. While we cannot review or discuss every possible scenario in this training, we will outline some general strategies for you to use with unclear situations.

Objectives

Participants will:

- » Identify strategies to resolve unclear situations

Exercise

- » Let's take a moment to reflect on what we have discussed. Using **Worksheet 6B: Unclear Situations**, take what we have gone through so far and think of persons that you have previously worked with or are working with currently or of a situation you imagine you may come across. Take a moment to write down several things that would make a case or situation unclear. For example, consider an individual who has a past history of suicidal crises, yet has denied suicidal ideation and, currently, has a strong support system.

- » Strategies to include for responding to situations that feel unclear:
 - Consult, internally or externally.
 - Refer, such as for further assessment, treatment, or assistance.
 - Develop a safety plan and contingency plan, including crisis resources and reducing access to lethal means if possible.
 - Involve support, mutually engaging the person's identified support network.
 - Follow up in a supportive and collaborative manner, using trauma-informed approaches and honoring safety.

Conclusion

Key Points

- » It is important to be cognizant of and identify linkages in suicide risk, trauma, and substance use.
- » While it may be complex and complicated to distinguish suicidal behavior and substance misuse, it is important to consider a holistic integrated treatment approach.
- » When situations seem less clear, strategies such as consultation, additional assessment, creating an expanded support network, and follow-up are particularly useful.

Module 7: Support the Family



Look for Hints

Ask Openly about
suicide

Validate Pain

Explore reasons to live
and plan to stay safe

H.O.P.E. saves lives

Module 7 Overview

+ Topics Covered


- » Common family reactions to a family member in crisis
- » Importance of maintaining a nonjudgmental stance
- » Ways to support the family across the spectrum of resource intensity



+ Learning Objectives

In Module 7, participants will:

- » Recognize the full range of normal reactions to a person at risk for suicide and the importance of maintaining a nonjudgmental stance
- » Identify interventions to support family members of a person struggling with suicidal thoughts and feelings



+ Materials

- » **Handout 7A: How to Talk about Suicide with a Loved One**



Introduction

Introduction

Now, we will be focusing on considerations to support the community, nation, and family that might be concerned about a person at risk for suicide or who have been impacted by a suicide loss. As we go into this section, let's keep in mind the importance of self-care, community care, leveraging all your resources, and staying in your role. You don't need to be all things to all people, and sometimes, it can actually be harmful to be so. While going through this information, think about how you have been doing this well in your role; others, including elders who have been supporting the community; and what can be learned from these strengths.

Common Family Reactions

Objectives

- » Recognize the full range of normal reactions to a person at risk for suicide and the importance of maintaining a nonjudgmental stance

Key Points

- » When a person you know and care about is thinking about suicide, it can evoke feelings of fear, anxiety, and helplessness. Often a family member's own need for that person to live can impact their ability to respond in a rational manner. Some family and friends may be able to work past this initial fear and tap into empathy, compassion, and concern.
- » Cultural and traditional beliefs and approaches to suicide may also impact family reactions. If the tribal tradition is to talk about death and suicide, the family may also do so in a compassionate way. If the traditional belief is to not talk about suicide, it is possible the family may avoid talking about it, feel more shame, or try to silence a person in the family who is struggling with suicidal thoughts. As you might remember from Module 2, strong family and cultural connections are protective factors against suicide. Therefore, it is important to consider how you as an advocate can strengthen the family while also being within the laws and ethical boundaries of your role.
- » There are some common thoughts that a concerned family member may have. These include thoughts such as:

- “Don’t they know I love them and would be devastated without them? How could they even think about this?”
- “They are always saying they want to kill themselves.”
- “Suicide is selfish.”
- “Suicide is cowardly.”
- “Suicide is a sin.”
- “You don’t mean that. You don’t really want to die.”
- “They’ve had so much going on. I hope I can support them through this.”

We are going to unpack these a little more:

- » Remember from our first section that a perception of being a burden is a common thought for the person in crisis. The tunnel vision of the suicidal mind can make the person think they are negatively impacting loved ones more in life than they would in death.
- » It can be invalidating and dismissive to think that a person who talks about suicide often is not going to act on it. The mind of the person in crisis can become used to thinking about suicide. Suicide can become the mind’s go-to solution or a rut it gets stuck in. Yet, this does not mean they won’t act on these thoughts at some point.
- » Saying that suicide is selfish can be a guilt- or shame-inducing response. If you think about crime victims, they may be feeling a lot of guilt and shame already connected to the trauma they have experienced. For persons who have lost someone to suicide, especially family members, this can be particularly hard. Thoughts such as “You know what we went through before. Why would you do this to us?” can surface. These can evoke anger and distance the supporter and the person struggling. A family can strengthen a victim’s connections while also refraining from statements that might distance a victim at risk.
- » Saying that suicide is a sin to someone struggling with suicidal thoughts could potentially increase alienation and hopelessness. It is not your job to change their belief, but you can help them to recognize that communicating this belief can be alienating to the person at risk. Connecting to beliefs such as compassion and love could help.
- » **Handout 7A: How to Talk about Suicide with a Loved One** provides a number of examples for how to talk about suicide with a loved one. This can be used as tool. It is best to meet the person where they are at, using their language. Consider what you would add or how you would change these.

Handout 7A: How to Talk about Suicide with a Loved One

There is more than one way to respond to a person struggling with a suicidal crisis. The following is a list of statements that may be helpful. It is essential to consider the individual, culture, and context and to use responses that are more natural to you. It is key to ask openly about suicide, connect with their pain, offer hope, and explore reasons to live while maintaining a collaborative relationship.

1. "I'm so glad you told me that you are thinking about suicide. I want you to be able to share whatever pain you are feeling."
2. "Help me understand what is going on that makes you want to die."
3. "How long have you been feeling bad/sad?"
4. "When did you start to lose hope that things could get better?"
5. "I care about you, and I would be so sad if I lost you to suicide."
6. "What can I do to help?"
7. "Are there things you have tried to do that didn't help?"
8. "What has been helpful to you during times when you have felt like you can't get through this?"
9. "Would you be willing to try some sources of help that you may not have tried before?"
10. "Have you thought about specific ways that you could kill yourself? What ways do you think are most likely?"
11. "I hear the intense pain you are in. I also hear that you would like to find a way through this pain. While I may not have all the answers, let's explore some ways to help you through this."
12. "You are such an important part of many peoples' lives, and losing you would be very painful and a deep loss for all who know you."



Considerations for support:

- » A victim who is already experiencing stress and trauma trying to help another person in crisis might struggle to know what to say. Problem-solving and providing specific examples can be very helpful. If it is the victim concerned about a family member, some education, coaching, and expanding the support system could be very helpful.
- » Family and friends may respond in a variety of different ways. It can help to find ways to tap into the more useful responses. Providing assistance to the support person may help them to be able to remain in a fully collaborative and supportive role.
- » Let's consider the variety of relationships and dynamics people have with those in crisis. Responses can vary depending on the relationship. For instance, a parent may feel guilty, like they have failed their child, if their child doesn't want to live. This can impact their response.
- » Think of a partner that may be suspected of interpersonal violence reporting that their spouse is thinking of suicide. At times, this "concern" can be used as a way to manipulate, control, or punish. This could be used to get the person in the hospital. It can also provide an opportunity to engage in other controlling behaviors, such as going through their phone or email or installing video cameras or monitoring devices in their car.
- » Also, consider if a person who is accused of inflicting harm or a perpetrator is making suicidal comments. These can still be serious suicidal comments. Yet, the victim is not in a position to support or respond and likely needs to be protected from this. Suicidal ideation, statements, and behaviors by the perpetrator could be triggering to the victim, or they could potentially cause the victim to be more concerned about the perpetrator than about themselves. This could compromise their safety. For instance, imagine someone has a protective order against their spouse, and they read a suicidal comment that the spouse made on social media. They might go to the spouse's house in the interest of helping that person remain safe from themselves. This could potentially jeopardize their own safety.
- » At the same time, what we know about persons who engage in violent, abusive, or harassing behaviors is that they may also have experienced trauma, adverse childhood experiences, and victimization. They can be at risk for suicide, as well. It is important to take suicidal statements and behaviors seriously while at the same time maintaining the safety of the victim. This is when a team approach is often best.

Supporting the Family

Introduction

We've been focusing on increasing our understanding and awareness of what the family or support persons might be experiencing. Understanding the complexities, background, and dynamics can help us maintain a nonjudgmental collaborative stance. In this next section, we will focus on ways you can support the family. Front and center is maintaining a collaborative, empathic relationship. At any time, you can establish a boundary and engage resources to support the family if the situation calls for this.

There can be a full range of reactions to a person at risk for suicide. This can be particularly challenging for family members, supporters, and others. While we can work to put ourselves in the shoes of another person, it is not possible to see fully and only from their perspective. It is also important to maintain the foundations of trauma-informed approaches and cultural humility in working with those impacted by suicide.

Objectives

- » Identify interventions to support children and adult family members of a person struggling with suicidal thoughts and feelings

Key Points

- » There are many ways to respond to a family member concerned about a person at risk for suicide. Be prepared ahead of time. Provide resources, make referrals, take action together when possible and appropriate, and engage the person's safety net. It may be helpful to consult with your team, and at times, you may need to take immediate action by initiating a crisis response.
 - You might respond directly to the person at risk, or you might only provide resources. It depends on the situation and your role. It is often useful to expand the safety net, provide crisis and treatment information, and consult. If you have a close relationship with the victim and it would not escalate their risk, you might consider reaching out directly. If the support person is concerned, the victim is in the process of a suicide attempt, or the safety of the victim is in question, you might initiate an emergency response. Similar to the resources available to you, the family member can call the National Suicide Prevention Lifeline or a mental health professional to seek consultation. There are also many resources listed on **Handout 4B: Resources**, your tribal community, and the C-SSRS, including community cards that list the questions of the screening tool. The cards can be easily accessed through the Columbia Lighthouse Project, printed, and given to a concerned support person.

- » If a child is concerned their parent is at risk for suicide, consider how to expand their support system, including traditional healers and medicine, to create a crisis response plan and help to identify a key adult who can support the parent.
- » You might assist the child in setting boundaries so they are not taking responsibility for the safety of their parent—both with suicide risk and protection from victimization. Helping the child to get support can be very useful.
- » It can be challenging for a spouse or partner to support a victim of trauma and through a suicidal crisis. Providing information about crisis support and treatment options and expanding their safety net can be useful responses. It can be helpful to problem solve together, educate the person that is concerned, and validate their limits by encouraging a safety net that is more than one person.
- » If a parent is concerned about a child at risk for suicide, consider how to expand their support system, provide crisis and emergency resources specific to youth, create a crisis response plan through emergency contacts, and help them think about how to meet their own needs. Set boundaries around your role and how you can assist them. Remind them who to contact for treatment and a mental health emergency. If the parent is having difficulty understanding the intensity of risk, provide support, educate, and engage a mental health professional or tribal leader, if appropriate.
- » It can be good to refrain from assuming that a parent is a “safe” or “helpful” presence. We recommend suggesting to the parent(s) that they should offer their child an opportunity to meet with a professional, and you can offer to be the go-between. Remember, you may not know the trauma history of this family.
- » **Reflection:** Take a moment to reflect on these questions:
 - Thinking about a suicidal crisis, when might a family member need to involve emergency crisis services for a person’s safety?
 - When might the family need to use a resource like a hospital?

Safer Suicide Care Within the Family

- » It may be helpful to prepare family and friends in how to support and recognize a crisis. These are key things to educate the family about:
 - Warning signs of suicide risk and how to respond.
 - If an attempt is in progress, call 911 for an emergency response.
 - If the victim has made a direct suicidal statement and cannot be reached or found to determine their safety, a welfare check may be needed.
 - If the person is not following their safety or treatment plan, a mental health crisis line or emergency mental health treatment may be needed.

Supporting Suicide Loss Survivors

- » At times, we may lose someone to suicide despite everyone's best efforts. This can contribute to deep hurt and pain. If someone you are supporting or the crime victim loses someone to suicide, connect to your community and resources that support you in healing. Suicide loss survivors can struggle with a range of feelings and reactions, including guilt, fixated thoughts on past interactions and their own behaviors, and questioning of reasons for the suicide death. You can listen without trying to have the answers. Support them in the way that works for your role, in a way that is collaborative and supportive. You might need to problem solve regarding specific tasks, such as who to contact or what needs to be addressed now. In other situations, it may be supportive to step back.
- » If a crime victim has died by suicide, your role may be particularly useful in helping to know how to handle an offender or an accused who may want to know, might start showing up, or might come to the funeral. If certain conditions exist, the offender may have rights to certain property. The family may need a lawyer, and you will likely need to use your team and consult. As this may be a challenging time, you may need to access your own support and professional assistance.

It is vital that all advocates be aware of the appropriate VOCA, VAWA, and/or FVPSA confidentiality requirements, depending on the funding source for their work. All of these funding organizations require that advocates have a signed, time-limited, written authorization for release of information from a victim who is served with funds from one of those funding streams before they can discuss what's going on with the victim, with the family, or with another third party.

- » How we respond to a suicide loss in a community can impact the potential ripple effect of suicide. Good support following a suicide loss can serve as suicide prevention and support healing, resiliency, and community connectedness.
 - Those impacted can be at higher risk for suicide themselves. Follow the best practices outlined in the safe messaging guidelines. It is also best to follow the postvention best practices described in the toolkits.
 - There are many resources for responding after a suicide loss, including:
 - A postvention toolkit for schools – <http://www.sprc.org/sites/default/files/resource-program/AfteraSuicideToolkitforSchools.pdf>
 - A toolkit for workplaces – <https://theactionalliance.org/sites/default/files/managers-guidebook-to-suicide-postvention-web.pdf>
 - Resources for safe messaging – <http://suicidepreventionmessaging.org>
- » Tap into the resilience, healing, and power of your community, including traditional ways and practices

- » If the crime victim is impacted by a suicide loss, there can be many complicating factors. Consider adding the intense grief of this type of loss to all of the experiences and impact of trauma. This can be overwhelming, can impact the victim's ability to function, and can stunt healing and recovery. The loss would be a precipitating event that might trigger a suicidal crisis or other crises for the victim. This would be a time to screen for suicide risk. In addition to identifying any risk, do the other things that you do so well, such as increasing contacts, connecting with support and treatment if needed, and using trauma-informed approaches. Additional complications can occur if the offender dies by suicide. This can have many possible reactions and impacts, including increased feelings of shame, guilt, blame, anger, and fear.
- » Losing a close support person to suicide can leave a victim feeling as if they don't have support or shouldn't burden others with their needs. They might feel guilty for having used the person for support. There can be many possible reactions that could complicate grief and healing in this situation.
- » It is important to keep in mind the many complicating factors for crime victims if they lose a person to suicide.

Conclusion

Key Points

- » We have outlined possible reactions of those who might be concerned a victim is at risk for suicide and ways to support them. We have also discussed ways to support the family, community, and victim if they are impacted by a suicide loss.
- » Knowing that you are one person, leverage your team and resources. Stay in your role, engage in self-care, and get support for yourself when needed.
- » Let's remember most people who attempt suicide go on to live without dying by suicide. The support makes a difference for those impacted, and you are not alone in this work.

Module 8: Consider the System



Look for Hints

Ask Openly about
suicide

Validate Pain

Explore reasons to live
and plan to stay safe

H.O.P.E. saves lives

Module 8 Overview

+ Topics Covered

- » Specific challenges and strengths of each crime victim setting
- » Strategies and considerations in developing suicide triage care policy

+ Learning Objectives

In Module 8, participants will:

- » Identify recommendations to enable settings to better support suicide prevention and care
- » Identify key champions within the organization and larger community to partner with for suicide prevention

+ Materials

Worksheet 8A: Communities and Systems

Introduction

Introduction

We have focused on supporting the victim, family, and community and your own self-care. Now we are moving on to discuss the communities in which each of you work. We encourage you to channel solution-focused brainstorming on this final section. It is okay to identify challenges, limits, and gaps. Yet, we can't stop there. We need to use our energies and collective wisdom to strengthen our communities and organizations. Trainings such as this can serve as an impetus for change.

Communities and Systems

Objectives

- » Identify recommendations to enable nations, communities, and organizations to support suicide prevention and care.

Key Points

- » We will focus on the first competency of developing recommendations to enable each setting to better support those at risk for suicide and each of you doing this work. We'll focus on identifying strengths, discussing possible strategies for improvement, and how to standardize processes.
- » **Reflection:** What are key strengths of your leadership? (This could be tribal leadership and leadership in the system in which you are working)
 - What is working well in suicide care in your organization or community?
 - What could be improved?
- » Now that you have considered some strengths of your community and organization, let's review some common challenges. While these challenges are not specific to your community or organization, as you read through them, think of which apply more to your specific area.
- » Often, leaders and systems may not have established clear standards, processes, or policies:
 - At times in systems, the "who to go to" person is identified. Yet, those on the ground aren't trained or provided the appropriate information to do this. For instance,

the “who to go to” person might be a crisis mental health professional, but the information for how to reach them, what to communicate, and options for their response may not be in the hands of the advocate when needed.

- Insufficient training can emphasize differences in individual responses. For instance, an untrained advocate might mistakenly judge that a victim is not at risk for suicide or miss hints.
- Poor documentation can occur when previous action steps, including safety planning, or previous identification of risk are not clearly documented. Imagine a survivor communicating signs of suicide risk, but the advocate responding in that specific moment doesn’t have the documentation that the person attempted suicide previously.
- Gaps in communication can refer to internal, external, and between-system communication.
 - Many systems do not consider resources for suicide loss until after suicide occurs.

There are several strategies to improve structure and consistency related to suicide prevention in crime victim advocate agencies:

- » Create standard training and protocols specific to roles. Be intentional to create this structure across the continuum of prevention, intervention, and postvention.
- » Clearly define roles and scope
 - Outline standard processes and procedures, including who to go to, who to inform and consult with, crisis policies, and standardized screening and triage, as possible.
- » Be specific about documentation within legal and ethical requirements. Yet, work to ensure that if many staff provide services to a person at risk, information is appropriately shared specifically, only the information that addresses risk and works to improve safety.
- » Using memorandums of understanding or memorandums of agreement to clearly outline partnerships.

Activity: Writing Exercise Focused on System Policies

Using **Worksheet 8A: Communities and Systems** focus on policies, opportunities for improvement, and what support is available to make such improvements.

Worksheet 8A: Communities and Systems

What type of policies do you know exist for your organization?

What do these policies communicate about your role in addressing and/or preventing suicide?

Where are the opportunities for improvement in these policies?

Who could help support you in seeking changes in your organization and community?



Identifying Champions

Introduction

Community and organizational changes typically don't happen overnight and by one person alone. Improvements occur more effectively and efficiently with engaged champions.

Objectives

- » Identify key champions within the organization and larger community to partner with for suicide prevention

Key Points

- » Focus on strengths, including strategies to leverage strengths and the power of champions.
- » Build on what works well in your communities and culture.
- » If you all do a great job supporting each other, consider what organizational supports and cultural changes made this happen and build on these approaches. An organization may have made progress in one area that can serve as momentum in another area.
- » A just culture is an organizational culture where the focus of reducing errors and preventing adverse events is on the system and not the individual. A just culture is a model of shared accountability, where feedback is sought at all levels, with no deference to rank or position, to constantly progress toward what is best for the client and safety. Errors are seen as learning opportunities to improve the system. A just, safety-oriented culture can be a foundation for suicide prevention with crime victims while taking the pressure off of the heroic efforts of you as an advocate. You have a system that is built to reduce gaps and support safety, healing, and recovery.

Reflection

- » **Identify:** Who are the champions in your organization or community? Who are the key leadership, staff, and other champions whom you can get on board?
- » Champions don't have to have a certain title. Yet, it is helpful to engage those who have some power and influence to change a system.
- » Engage those who are passionate about the issue or improvement you are seeking. Also, work to engage key leaders who have the power to impact change. Use both stories and

data, when appropriate, to leverage and engage champions. When possible, work to get those with expertise or lived experience on board. People with lived experience are those who have been through a crisis and can use their experiences to inform systemic and community changes to better support others who experience suicidal crises.

- » If it feels too challenging to seek systems change and still care for yourself and others, consider how to have a voice in a way that works best for you and your system.

Conclusion

Key Points

- » Suicide prevention requires an organizational commitment and a systematic approach.
- » The work of supporting the safety and recovery of those we are supporting must be a team effort and not on the heroic efforts of individual advocates.
- » Organizations need to review training, policies, procedures, and referral networks to develop a comprehensive approach to suicide prevention.
- » Leveraging the energy and passion of champions can create momentum for systemic change.

Training Wrap-Up

We have come to the end of our training program. To pull it all together, we've discussed the specific suicide risk data that crime victims face and have discussed that certain communities are at higher risk than others. We've also discussed the following topics:

1. The role of the crime victim advocate
2. The kinds of hints to listen for that may signal suicidal ideation and risk and how to listen and openly communicate that you are listening and unafraid of discussing suicide
3. The best ways to address suicidal statements and to screen for suicide risk
4. The importance of addressing suicide directly and openly after indirect questions
5. Strategies to validate pain, explore reasons for living, and establish a safety plan
6. Important issues that may co-exist with suicide risk, including trauma and substance use, and the ways that co-occurring issues can exacerbate or inform suicide risk

7. The best ways to help family members and close supports who may be within the suicidal person's social network
8. The importance of taking care of our own mental health and well-being through intentional self-care
9. The importance of recognizing and working to improve the systems we each work within, so that our advocacy can be most effective and so that we can be the most effective.

This training, developed by Education Development Center with support and in close collaboration with the Office for Victims of Crime, is just one foundational step in the learning available to you. Access resources, videos, and additional information about the H.O.P.E. Suicide Prevention Training on the EDC website: <https://www.edc.org/hopeforadvocates>.

There are resources available, including training and technical assistance, through the [Office for Victims of Crime](#).

[Education Development Center](#), including its [Suicide Prevention Resource Center](#) and its [Zero Suicide Institute](#), is available for consultation and questions. We offer a large array of publicly available webinars, resources, and training that you and your organization are welcome to take advantage of. If your organization is interested in having a H.O.P.E Trainer organize a training for your staff, please contact Dr. Heidi Kar at hkar@edc.org.

In addition to these resources, the [Indian Health Service](#) and the [Substance Abuse and Mental Health Services Administration](#) have many resources regarding substance misuse, suicide prevention, trauma-informed practices, and more.

The [National Action Alliance for Suicide Prevention](#) is the nation's public-private partnership focused on suicide prevention.

The [National Suicide Prevention Lifeline](#) (1-800-273-8255) provides crisis services for people in distress, their loved ones, and best practices for professionals.

References

- American Counseling Association. (n.d.). *Fact sheet #9: Vicarious trauma*. https://www.counseling.org/docs/trauma-disaster/fact-sheet-9---vicarious-trauma.pdf?sfvrsn=f0f03a27_2
- American Foundation for Suicide Prevention. (2020). *Risk factors and warning signs*. Retrieved June 22, 2020, from <https://afsp.org/risk-factors-and-warning-signs>
- American Psychological Association. (2020). *Trauma*. Retrieved June 23, 2020, from <https://www.apa.org/topics/trauma/>
- Centers for Disease Control and Prevention. (2020). *Wide Ranging Online Data for Epidemiological Research (WONDER), Cause of death files, 1999-2018* [Data file]. <http://wonder.cdc.gov/ucd-icd10.html>
- Chu, C., Buchman-Schmitt, J. M., Stanley, I. H., Hom, M. A., Tucker, R. P., Hagan, C. R., Rogers, M. L., Podlogar, M. C., Chiurliza, B., Ringer, F. B., Michaels, M. S., Patros, C., & Joiner, T. E. (2017). The interpersonal theory of suicide: A systematic review and meta-analysis of a decade of cross-national research. *Psychological Bulletin*, 143(12), 1313–1345. <https://doi.org/10.1037/bul0000123>
- Columbia Lighthouse Project. (2016). *The Columbia Protocol for healthcare and other community settings*. Retrieved June 22, 2020, from <https://cssrs.columbia.edu/the-columbia-scale-c-ssrs/cssrs-for-communities-and-healthcare/>
- Columbia Lighthouse Project. (2018). *Columbia-Suicide Severity Rating Scale: Screen with triage points for victim advocates*. <https://cssrs.columbia.edu/documents/screener-recent-with-triage-for-clinicians-advocates/c-ssrs-screener-with-triage-points-for-victim-advocates-2018/>
- Doupnik, S. K., Rudd, B., Schmutte, T., Worsley, D., Bowden, C. F., McCarthy, E., Eggan, E., Bridge, J. A., & Marcus, S. C. (2020). Association of suicide prevention interventions with subsequent suicide attempts, linkage to follow-up care, and depression symptoms for acute care settings: A systematic review and meta-analysis. *JAMA Psychiatry*. <https://doi.org/10.1001/jamapsychiatry.2020.1586>
- Ellis, T. E. (2006). *Cognition and suicide: Theory, research, and therapy*. American Psychological Association.
- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., & Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14(4), 245–258. [https://doi.org/10.1016/s0749-3797\(98\)00017-8](https://doi.org/10.1016/s0749-3797(98)00017-8)
- Harris, R. (2009). *ACT made simple: An easy-to-read primer on acceptance and commitment therapy*. New Harbinger Publications.
- Institute for Quality and Efficiency in Health Care. (2017, January 12). *Depression: What is burnout?* <https://www.ncbi.nlm.nih.gov/books/NBK279286/>

- Leavitt, R. A., Ertl, A., Sheats, K., Petrosky, E., Ivey-Stephenson, A., & Fowler, K. A. (2018). Suicides among American Indian/Alaska Natives — National Violent Death Reporting System, 18 states, 2003–2014. *Morbidity and Mortality Weekly Report*, 67(8), 237–242. <https://doi.org/10.15585/mmwr.mm6708a1>
- Levi-Belz, Y., Gvion, Y., & Apter, A. (2019). Editorial: The psychology of suicide: From research understandings to intervention and treatment. *Frontiers in Psychiatry*, 10(214). <https://doi.org/10.3389/fpsy.2019.00214>
- Millner, A. J., Lee, M. D., & Nock, M. K. (2016). Describing and measuring the pathway to suicide attempts: A preliminary study. *Suicide and Life-Threatening Behavior*, 47(3), 353–369. <https://doi.org/10.1111/sltb.12284>
- Morse, G., Salyers, M. P., Rollins, A. L., Monroe-Devita, M., & Pfahler, C. (2012). Burnout in mental health services: A review of the problem and its remediation. *Administration and Policy in Mental Health and Mental Health Services Research*, 39(5), 341–352. <https://doi.org/10.1007/s10488-011-0352-1>
- Now Matters Now. (2018). *Free training and resources*. Retrieved June 26, 2020, from <https://www.nowmattersnow.org/get-involved>
- Office for Victims of Crime. (n.d.a). *Vicarious Trauma Toolkit: About the Toolkit*. Retrieved June 22, 2020, from <https://vtt.ovc.ojp.gov/about-the-toolkit>
- Office for Victims of Crime. (n.d.b). *Vicarious Trauma Toolkit: Introduction*. Retrieved June 22, 2020, from <https://ovc.ojp.gov/program/vtt/introduction>
- Office for Victims of Crime. (n.d.c). *Vicarious Trauma Toolkit: What is vicarious trauma?* Retrieved June 22, 2020, from <https://vtt.ovc.ojp.gov/what-is-vicarious-trauma>
- Oltmanns, T. F. (2017). *Common traits*. Florida Initiative for Suicide Prevention. Retrieved June 25, 2020, from <https://fisponline.org/programs/education/common-traits/>
- Oquendo, M. A., & Volkow, N. D. (2018). Suicide: A silent contributor to opioid-overdose deaths. *New England Journal of Medicine*, 378(17), 1567–1569. <https://doi.org/10.1056/NEJMp1801417>
- Pompili, M., Serafini, G., Innamorati, M., Dominici, G., Ferracuti, S., Kotzalidis, G. D., & Sher, L. (2010). Suicidal behavior and alcohol abuse. *International Journal of Environmental Research and Public Health*, 7(4), 1392–1431. <https://doi.org/10.3390/ijerph7041392>
- Ports, K. A., Merrick, M. T., Stone, D. M., Wilkins, N. J., Reed, J., Ebin, J., & Ford, D. C. (2017). Adverse childhood experiences and suicide risk: Toward comprehensive prevention. *American Journal of Preventive Medicine*, 53(3), 400–403. <https://doi.org/10.1016/j.amepre.2017.03.015>
- Rosay, A. B. (2016, June 1). *Violence against American Indian and Alaska Native women and men*. National Institute of Justice. <https://nij.ojp.gov/topics/articles/violence-against-american-indian-and-alaska-native-women-and-men>
- Sachs-Ericsson, N. J., Rushing, N. C., Stanley, I. H., & Sheffler, J. (2015). In my end is my beginning: Developmental trajectories of adverse childhood experiences to late-life suicide. *Aging & Mental Health*, 20(2), 139–165. <https://doi.org/10.1080/13607863.2015.1063107>

Schauben, L. J., & Frazier, P. A. (1995). Vicarious trauma: The effects on female counselors of working with sexual violence survivors. *Psychology of Women Quarterly*, 19(1), 49–64. <https://doi.org/10.1111/j.1471-6402.1995.tb00278.x>

Shneidman, E. S. (1998). *The suicidal mind*. Oxford University Press.

Spencer-Thomas, S. (2017, July). *Understanding the suicidal mind*. International Risk Management Institute. Retrieved June 25, 2020, from <https://www.irmi.com/articles/expert-commentary/understanding-the-suicidal-mind>

Stanford SPARQ. (n.d.). *Caring letters prevent suicide*. Stanford University. Retrieved June 22, 2020, from <https://sparq.stanford.edu/solutions/caring-letters-prevent-suicide>

Stanley, B., & Brown, G. K. (2008). *Patient safety plan template*. Retrieved June 22, 2020, from <http://www.sprc.org/resources-programs/patient-safety-plan-template>

Stanley, B., & Brown, G. K. (2012). Safety planning intervention: A brief intervention to mitigate suicide risk. *Cognitive and Behavioral Practice*, 19(2), 256–264. <https://doi.org/10.1016/j.cbpra.2011.01.001>

Stanley, B., & Brown, G. K. (2009). *Safety planning guide: A quick guide for clinicians*. <https://www.sprc.org/resources-programs/safety-planning-guide-quick-guide-clinicians>

Substance Abuse and Mental Health Services Administration. (2014). *SAMHSA's concept of trauma and guidance for a trauma-informed approach*. https://ncsacw.samhsa.gov/userfiles/files/SAMHSA_Trauma.pdf

Substance Abuse and Mental Health Services Administration. (2015). *In brief: Substance use and suicide: A nexus requiring a public health approach*. <https://store.samhsa.gov/sites/default/files/d7/priv/sma16-4935.pdf>

Substance Abuse and Mental Health Services Administration. (2018, June). *The role of adverse childhood experiences in substance misuse and related behavioral health problems*. <https://mnprc.org/wp-content/uploads/2019/01/aces-behavioral-health-problems.pdf>

Suicide Prevention Resource Center. (n.d.). *Warning signs for suicide*. Retrieved June 22, 2020, from <http://www.sprc.org/about-suicide/warning-signs>

Suicide Prevention Resource Center. (2020a). *Risk and protective factors*. Retrieved June 22, 2020, from <https://www.sprc.org/about-suicide/risk-protective-factors>

Suicide Prevention Resource Center. (2020b). *Suicidal thoughts and suicide attempts*. Retrieved June 24, 2020, from <http://www.sprc.org/scope/attempts>

University of Toronto. (2016). *Link found between witnessing parental domestic violence during childhood and attempted suicide: One in six adults who were exposed to chronic parental domestic violence during their childhood has attempted suicide*. ScienceDaily. Retrieved June 25, 2020 from www.sciencedaily.com/releases/2016/06/160609115306.htm

United States Department of Veterans Affairs. (2019, October). *Suicide and PTSD*. https://www.ptsd.va.gov/understand/related/suicide_ptsd.asp

United States Office of the Surgeon General & National Action Alliance for Suicide Prevention. (2012) *2012 National strategy for suicide prevention: Goals and objectives for action: A report of the U.S. Surgeon General and of the National Action Alliance for Suicide Prevention*. United States Department of Health & Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK109910/>

We R Native. (n.d.). *Honor LGBTQ2S people*. Retrieved June 22, 2020, from <https://www.wernative.org/blog-posts/honor-lgbtq2s-people>

Zero Suicide Institute. (2019). *Best and promising practices for the implementation of Zero Suicide in Indian Country*. http://zerosuicide.edc.org/toolkit/indian-country#quicktabs-native_american=5

Zero Suicide Institute. (2020a). *Zero Suicide toolkit: Transition*. Retrieved June 22, 2020, from <http://zerosuicide.edc.org/toolkit/transition>

Zero Suicide Institute. (2020b). *Zero Suicide toolkit: Treat*. Retrieved June 23, 2020, from <http://zerosuicide.edc.org/toolkit/treat>



www.edc.org

EDC 43 Foundry Avenue Waltham, MA 02453

E-mail: contact@edc.org

Phone: 617-969-7100

Boston | Chicago | New York | Washington, D.C.

Education Development Center (EDC) is a global nonprofit that advances lasting solutions to improve education, promote health, and expand economic opportunity. Since 1958, we have been a leader in designing, implementing, and evaluating powerful and innovative programs in more than 80 countries around the world.